

# Stream Function Formulation of Surface Stokes Equations

Arnold Reusken\*

Institut für Geometrie und Praktische Mathematik  
Templergraben 55, 52062 Aachen, Germany

---

\* IGPM, RWTH Aachen University, Templergraben 55, D-52062 Aachen, Germany (reusken@igpm.rwth-aachen.de)

# STREAM FUNCTION FORMULATION OF SURFACE STOKES EQUATIONS

ARNOLD REUSKEN\*

**Abstract.** In this paper we present a derivation of the surface Helmholtz decomposition, discuss its relation to the surface Hodge decomposition, and derive a well-posed stream function formulation of a class of surface Stokes problems. We consider a  $C^2$  connected (not necessarily simply connected) oriented hypersurface  $\Gamma \subset \mathbb{R}^3$  without boundary. The surface gradient, divergence, curl and Laplace operators are defined in terms of the standard differential operators of the ambient Euclidean space  $\mathbb{R}^3$ . These representations are very convenient for the implementation of numerical methods for surface partial differential equations. We introduce surface  $\mathbf{H}(\operatorname{div}_\Gamma)$  and  $\mathbf{H}(\operatorname{curl}_\Gamma)$  spaces and derive useful properties of these spaces. A main result of the paper is the derivation of the Helmholtz decomposition, in terms of these surface differential operators, based on elementary differential calculus. As a corollary of this decomposition we obtain that for a simply connected surface, to every tangential divergence free velocity field there corresponds a unique scalar stream function. Using this result the variational form of the surface Stokes equation can be reformulated as a well-posed variational formulation of a fourth order equation for the stream function. The latter can be rewritten as two coupled second order equations, which form the basis for a finite element discretization. A particular finite element method is explained and results of a numerical experiment with this method are presented.

**Key words.** surface Stokes, surface Helmholtz decomposition, stream function formulation

**1. Introduction.** In the literature on modeling of emulsions, foams or biological membranes mathematical models describing fluidic surfaces or fluidic interfaces occur; cf., e.g., [40, 41, 4, 7, 34, 33]. Typically such models consist of surface (Navier-)Stokes equations. These equations are also studied as an interesting mathematical problem in its own right in, e.g., [14, 44, 43, 3, 26, 2, 23]. Recently there has been a strong increase in research on numerical simulation methods for surface (Navier-)Stokes equations, e.g., [29, 5, 36, 35, 37, 16, 22, 30]. By far most of these and other papers on numerical methods for surface flow problems treat the (Navier-)Stokes equations in the primitive velocity and pressure variables. In the paper [29] the Navier-Stokes equations on a stationary smooth closed surface in *stream function* formulation are treated. We are not aware of any other literature in which surface (Navier-)Stokes equations in stream function formulation are studied.

In Euclidean space, the stream function formulation of (Navier-)Stokes is well-known and thoroughly studied, e.g., [17, 32] and the references therein. In numerical simulations of three-dimensional problems this formulation is not often used due to substantial disadvantages. For two-dimensional problems this formulation reduces to a fourth order biharmonic equation for the *scalar* stream function. This formulation has been used in numerical simulations, although it has certain disadvantages related to boundary conditions and regularity ([17, 32]).

In the fields of applications mentioned above, one often deals with smooth simply connected surfaces without boundary. In such a setting there usually are no difficulties related to regularity or boundary conditions and the stream function formulation may be a very attractive alternative to the formulation in primitive variables, as already indicated in [29]. This is the main motivation for the study presented in this paper. We present a detailed analysis of the stream function formulation for a certain class of surface Stokes equations. It is clear that such a stream function formulation should

---

\*Institut für Geometrie und Praktische Mathematik, RWTH-Aachen University, D-52056 Aachen, Germany (reusken@igpm.rwth-aachen.de).

be based on a Helmholtz decomposition of  $L^2(\Gamma)$  (where  $\Gamma$  denotes the surface). This decomposition can be interpreted as a variant of the Hodge-decomposition from the field of differential forms. It turns out that for this surface Helmholtz decomposition and the corresponding stream function formulation of the Stokes problem only some partial results are available in the literature. For example, in [8] a Helmholtz decomposition for the case that  $\Gamma$  is a simply connected Lipschitz polyhedron is studied and in [29] a stream function formulation is derived in the setting of differential forms.

In this paper we present a complete derivation of the surface Helmholtz decomposition, discuss its relation to the surface Hodge decomposition, and derive a well-posed stream function formulation of a class of surface Stokes problems. We consider a  $C^2$  connected (not necessarily simply connected) oriented hypersurface  $\Gamma \subset \mathbb{R}^3$  without boundary. We introduce the natural surface gradient, divergence, curl and Laplace operators, represented in terms of the standard differential operators of the ambient Euclidean space  $\mathbb{R}^3$ . These representations, which may differ from the (intrinsic) ones used in differential geometry, are very convenient for the implementation of numerical methods for surface PDEs. Similar representations for surface differential operators are also used in e.g., [21, 20, 16, 30, 37]. We introduce suitable surface  $\mathbf{H}(\operatorname{div}_\Gamma)$  and  $\mathbf{H}(\operatorname{curl}_\Gamma)$  spaces and derive useful properties of these spaces. A main result of the paper is the derivation of the Helmholtz decomposition (Theorem 4.2), in terms of these surface differential operators, based on elementary differential calculus. In particular, we do not use the calculus of differential forms. However, we do point out the relation between the Helmholtz decomposition and a Hodge decomposition known from the field of differential forms. As a corollary of this Helmholtz decomposition we obtain that for a simply connected surface, to every tangential divergence free velocity field there corresponds a unique scalar stream function. Using this result the variational form of the Stokes equation can be reformulated as a well-posed variational formulation of a fourth order equation for the stream function. The latter can be rewritten as two coupled second order equations, which form the basis for a finite element discretization.

The remainder of the paper is organized as follows. In Section 2 we introduce surface differential operators and derive useful relations between these operators. Surface Sobolev spaces, in particular  $\mathbf{H}(\operatorname{div}_\Gamma)$  and  $\mathbf{H}(\operatorname{curl}_\Gamma)$ , are introduced in Section 3 and some basic properties of these spaces are derived. In Section 4 the surface Helmholtz decomposition is presented and a few corollaries, e.g., a Friedrichs type inequality for tangential velocity vectors, are treated. Furthermore it is explained how the Helmholtz decomposition relates to a certain Hodge decomposition. In Section 5, for the case of a simply connected surface  $\Gamma$ , a class of surface Stokes problems is discussed and a reformulation in terms of a well-posed problem for the stream function is treated. Finally, in Section 6 we present results of a numerical experiment for a finite element discretization of the stream function formulation.

**2. Surface differential operators.** In this section we introduce surface differential operators for smooth ( $C^1(\Gamma)$ ) functions and derive properties for these operators. We consider a sufficiently smooth closed connected compact surface  $\Gamma \subset \mathbb{R}^3$ . In this paper we do not try to derive results under minimal smoothness conditions for the surface  $\Gamma$ . We introduce the following assumption, which is sufficient (but not necessary) for our analysis.

ASSUMPTION 2.1. *In the remainder of the paper we assume that  $\Gamma$  is a  $C^2$  connected compact oriented hypersurface in  $\mathbb{R}^3$  without boundary.*

There are different ways for introducing tangential and covariant derivatives for scalar,

vector or matrix valued functions defined on  $\Gamma$ . In differential geometry one uses the notion of covariant derivative, which is intrinsic for a Riemannian surface, i.e., one does not use the embedding of a surface in an ambient space [11, 24]. A related more general concept of derivatives is introduced in exterior calculus via the exterior derivative of differential forms, cf. [1]. We will comment further on this in Section 4.1. In this paper we represent differential operators on  $\Gamma$  by making explicit use of the embedding Eulerian space  $\mathbb{R}^3$ . The motivation for this comes from numerical analysis. In recent papers on numerical methods for surface PDEs it has been shown that the formulation of surface PDEs in terms of these differential operators is very convenient for numerical simulation, e.g., the review paper [13] for scalar surface PDEs and [21, 36, 37, 29, 30, 38] for surface (Navier-)Stokes equations. In particular in the setting of surface Stokes equations the  $\nabla_\Gamma$ ,  $\text{div}_\Gamma$  and  $\mathbf{curl}_\Gamma$  differential operators introduced below play a key role. We summarize some basic properties of these operators and derive a relation ((2.14) below) that relates the surface  $\mathbf{curl}_\Gamma$  differential operator to surface vector Laplacians, cf. Remark 2.1. The analysis is elementary, using basic tensor analysis.

The outward pointing unit normal and the signed distance function are denoted by  $\mathbf{n}$  and  $d$ , respectively. On a sufficiently small neighborhood  $U$  of  $\Gamma$  the closest point projection is given by  $\mathbf{p}(x) = x - d(x)\mathbf{n}(x)$ . We also use the orthogonal projection  $\mathbf{P}(x) = \mathbf{I} - \mathbf{n}(x)\mathbf{n}(x)^T$ ,  $x \in \Gamma$ . The tangential derivative of a scalar function  $\phi \in C^1(\Gamma)$  and of a vector function  $\mathbf{u} \in C^1(\Gamma)^3$  are, for  $x \in \Gamma$ , defined by

$$\nabla_\Gamma \phi(x) = \nabla(\phi \circ \mathbf{p})(x) = \mathbf{P}(x)\nabla\phi^e(x), \quad (2.1)$$

$$\begin{aligned} \nabla_\Gamma \mathbf{u}(x) &= \left( \frac{\partial(\mathbf{u} \circ \mathbf{p})(x)}{\partial x_1} \quad \frac{\partial(\mathbf{u} \circ \mathbf{p})(x)}{\partial x_2} \quad \frac{\partial(\mathbf{u} \circ \mathbf{p})(x)}{\partial x_2} \right) \mathbf{P}(x) \\ &= \mathbf{P}(x)\nabla\mathbf{u}^e(x)\mathbf{P}(x), \end{aligned} \quad (2.2)$$

where  $\phi^e$ ,  $\mathbf{u}^e$  denote some smooth extension of  $\phi$  and  $\mathbf{u}$  on the neighborhood  $U$ , and  $\nabla\mathbf{u}^e$  is the Jacobian,  $(\nabla\mathbf{u}^e)_{i,j} = \frac{\partial u_i^e}{\partial x_j}$ ,  $1 \leq i, j \leq 3$ . In the remainder we delete the argument  $x \in \Gamma$ . If the vector function  $\mathbf{u}$  is tangential, i.e.,  $\mathbf{n} \cdot \mathbf{u} = 0$  on  $\Gamma$ , then  $\nabla_\Gamma \mathbf{u}$  coincides with the covariant derivative. We also need the tangential divergence operators corresponding to  $\nabla_\Gamma$ . In analogy to the definitions used for vector and matrix valued functions in Euclidean space  $\mathbb{R}^3$  we introduce

$$\text{div}_\Gamma \mathbf{u} := \text{tr}(\nabla_\Gamma \mathbf{u}), \quad \text{div}_\Gamma A := \begin{pmatrix} \text{div}_\Gamma(e_1^T A) \\ \text{div}_\Gamma(e_2^T A) \\ \text{div}_\Gamma(e_3^T A) \end{pmatrix}, \quad A \in C^1(\Gamma)^{3 \times 3}, \quad (2.3)$$

where  $e_i$ ,  $i = 1, 2, 3$ , are the standard basis vectors in  $\mathbb{R}^3$ . We recall well-known partial integration identities. For this we introduce the space of *tangential* vector functions  $C_t^m(\Gamma)^3 := \{\mathbf{u} \in C^m(\Gamma)^3 \mid \mathbf{n} \cdot \mathbf{u} = 0 \text{ on } \Gamma\}$ . The following relations hold:

$$\int_\Gamma \text{div}_\Gamma \mathbf{u} \phi \, ds = - \int_\Gamma \mathbf{u} \cdot \nabla_\Gamma \phi \, ds \quad \text{for all } \mathbf{u} \in C_t^1(\Gamma)^3, \phi \in C^1(\Gamma), \quad (2.4)$$

$$\begin{aligned} &\int_\Gamma (\text{div}_\Gamma A) \cdot \mathbf{u} \, ds \\ &= - \int_\Gamma \text{tr}(A^T \nabla_\Gamma \mathbf{u}) \, ds \quad \text{for all } \mathbf{u} \in C^1(\Gamma)^3, A \in C^1(\Gamma)^{3 \times 3} \text{ with } \mathbf{P}A\mathbf{P} = A. \end{aligned} \quad (2.5)$$

The relation (2.4) can be found at many places in the literature, e.g. [13]. The result in (2.5) can easily be derived using a componentwise application of (2.4). Hence the

$\nabla_\Gamma$  and  $\text{div}_\Gamma$  operator have the usual relation  $\nabla_\Gamma = -\text{div}_\Gamma^T$  in the sense (2.4), (2.5). We also need an appropriate surface curl operator. In analogy to the 2D curl operator  $\text{curl}_{2D} := (\nabla \times \mathbf{u}) \cdot \mathbf{e}_3$  it is given by

$$\text{curl}_\Gamma \mathbf{u} := (\nabla_\Gamma \times \mathbf{u}^e) \cdot \mathbf{n}, \quad \mathbf{u} \in C^1(\Gamma)^3. \quad (2.6)$$

The following useful identity holds; a proof is given in the Appendix:

$$\text{curl}_\Gamma \mathbf{u} = \text{div}_\Gamma(\mathbf{u} \times \mathbf{n}), \quad \mathbf{u} \in C^1(\Gamma)^3. \quad (2.7)$$

As adjoint of this surface curl operator we have the vector-curl operator defined by

$$\mathbf{curl}_\Gamma \phi := \mathbf{n} \times \nabla_\Gamma \phi, \quad \phi \in C^1(\Gamma). \quad (2.8)$$

Using (2.4) and the vector product rule  $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = (\mathbf{c} \times \mathbf{a}) \cdot \mathbf{b}$  we get

$$\begin{aligned} \int_\Gamma \text{curl}_\Gamma \mathbf{u} \phi \, ds &= \int_\Gamma \text{div}_\Gamma(\mathbf{u} \times \mathbf{n}) \phi \, ds = - \int_\Gamma (\mathbf{u} \times \mathbf{n}) \cdot \nabla_\Gamma \phi \, ds \\ &= - \int_\Gamma (\mathbf{n} \times \nabla_\Gamma \phi) \cdot \mathbf{u} \, ds = - \int_\Gamma \mathbf{u} \cdot \mathbf{curl}_\Gamma \phi \, ds. \end{aligned}$$

Hence,

$$\int_\Gamma \text{curl}_\Gamma \mathbf{u} \phi \, ds = - \int_\Gamma \mathbf{u} \cdot \mathbf{curl}_\Gamma \phi \, ds, \quad \text{for } \mathbf{u} \in C_t^1(\Gamma)^3, \phi \in C^1(\Gamma), \quad (2.9)$$

and thus indeed  $\text{curl}_\Gamma = -\mathbf{curl}_\Gamma^T$  holds.

In the following lemma we collect some relations. The relations (i)-(iii) are elementary. The result in (iv), however, requires a longer analysis. We comment on the result (iv) in Remark 2.1

LEMMA 2.1. *The following relations hold on  $\Gamma$ , for all  $\phi \in C^2(\Gamma)$ ,  $\mathbf{u} \in C_t^2(\Gamma)^3$ , with  $K = K(x)$  the Gauss curvature on  $\Gamma$ :*

$$(i) \quad \text{div}_\Gamma(\mathbf{curl}_\Gamma \phi) = 0 \quad (2.10)$$

$$(ii) \quad \text{curl}_\Gamma(\nabla_\Gamma \phi) = 0 \quad (2.11)$$

$$(iii) \quad \text{curl}_\Gamma(\mathbf{curl}_\Gamma \phi) = \text{div}_\Gamma(\nabla_\Gamma \phi) = \Delta_\Gamma \phi \quad (2.12)$$

$$(iv) \quad \mathbf{curl}_\Gamma(\text{curl}_\Gamma \mathbf{u}) = \mathbf{P} \text{div}_\Gamma(\nabla_\Gamma \mathbf{u} - \nabla_\Gamma \mathbf{u}^T) \quad (2.13)$$

$$= \mathbf{P} \text{div}_\Gamma(\nabla_\Gamma \mathbf{u}) - \nabla_\Gamma(\text{div}_\Gamma \mathbf{u}) - K \mathbf{u}. \quad (2.14)$$

*Proof.* From the definitions it follows that  $\text{div}_\Gamma(\mathbf{curl}_\Gamma \phi) = \text{div}_\Gamma(\mathbf{n} \times \nabla_\Gamma \phi) = -\text{curl}_\Gamma(\nabla_\Gamma \phi)$ . In the Appendix we derive

$$\text{div}_\Gamma(\mathbf{n} \times \nabla_\Gamma \phi) = 0. \quad (2.15)$$

From this the results in (2.10), (2.11) follow. The result in (2.12) follows from elementary properties of the vector product:

$$\text{curl}_\Gamma(\mathbf{curl}_\Gamma \phi) = \text{curl}_\Gamma(\mathbf{n} \times \nabla_\Gamma \phi) = \text{div}_\Gamma((\mathbf{n} \times \nabla_\Gamma \phi) \times \mathbf{n}) = \text{div}_\Gamma(\nabla_\Gamma \phi) = \Delta_\Gamma \phi.$$

The proof of (2.13) requires a longer tedious, but elementary, derivation that is given in the Appendix. In [22], Lemma 2.1, the identity  $\mathbf{P} \text{div}_\Gamma(\nabla_\Gamma \mathbf{u}^T) = \nabla_\Gamma(\text{div}_\Gamma \mathbf{u}) + K \mathbf{u}$

is derived. This yields the result (2.14).  $\square$

Note that in (2.13), (2.14) we use the surface divergence applied to a matrix. As a simple consequence of (2.13)-(2.14) we formulate the following identity that will be used in the derivation of the stream formulation of the surface Stokes problem. If  $\mathbf{u} \in C_t^2(\Gamma)^3$  satisfies  $\operatorname{div}_\Gamma \mathbf{u} = 0$  then the relation

$$\mathbf{P} \operatorname{div}_\Gamma (\nabla_\Gamma \mathbf{u} + \nabla_\Gamma \mathbf{u}^T) = \mathbf{curl}_\Gamma (\operatorname{curl}_\Gamma \mathbf{u}) + 2K\mathbf{u} \quad (2.16)$$

holds.

REMARK 2.1. The relations (2.13), (2.14) are key identities for the reformulation of Stokes equations in stream function formulation. We are not aware of a rigorous proof of these relations in the literature. In [29] similar relations for surface curl operators defined via  $k$ -forms are discussed. Note that in our setting we define all surface differential operators through the Euclidean differential operators in the embedding space  $\mathbb{R}^3$  (avoiding  $k$ -forms) and the proofs of (2.13), (2.14) are based on elementary tensor analysis. We briefly discuss how the identities (2.13), (2.14) are related to well-known ones in Euclidean space  $\mathbb{R}^2$ . If  $\Gamma \subset \mathbb{R}^2$  the definitions (2.6), (2.8) yield for  $\mathbf{u} = (u_1, u_2)$ :

$$\operatorname{curl}_{2D} \mathbf{u} = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y}, \quad \mathbf{curl}_{2D} \phi = \left( -\frac{\partial \phi}{\partial y}, \frac{\partial \phi}{\partial x} \right)^T.$$

Note that these are the standard curl definitions, apart from a sign change in  $\mathbf{curl}_{2D}$ . A basic identity found at many places in the literature ([17]) is the following:

$$\mathbf{curl}_{2D} (\operatorname{curl}_{2D} \mathbf{u}) = \Delta \mathbf{u} - \nabla (\operatorname{div} \mathbf{u}). \quad (2.17)$$

Using  $\Delta \mathbf{u} = \operatorname{div} (\nabla \mathbf{u})$ ,  $\nabla \operatorname{div} \mathbf{u} = \operatorname{div} (\nabla \mathbf{u}^T)$  this can be rewritten as

$$\mathbf{curl}_{2D} (\operatorname{curl}_{2D} \mathbf{u}) = \operatorname{div} (\nabla \mathbf{u} - \nabla \mathbf{u}^T). \quad (2.18)$$

We see that the latter relation has exactly the same form as the surface identity (2.13), whereas in the generalization of (2.17) to its surface variant (2.14) an additional curvature term  $K\mathbf{u}$  enters. Finally we note that the relation (2.14) is closely related to the so-called Weitzenböck identity from differential geometry, cf. (4.11).

We finally recall two Stokes type identities on a connected Lipschitz subdomain  $\gamma \subset \Gamma$ . The outward pointing unit normal to  $\partial\gamma$  that is tangential to  $\Gamma$  is denoted by  $\boldsymbol{\nu}$ . The induced vector tangential to both  $\partial\gamma$  and  $\Gamma$  is denoted by  $\boldsymbol{\tau} := \mathbf{n} \times \boldsymbol{\nu}$ . The following Stokes relations hold:

$$\int_\gamma \operatorname{div}_\Gamma \mathbf{u} \, ds = \int_{\partial\gamma} \mathbf{u} \cdot \boldsymbol{\nu} \, ds, \quad \text{for } \mathbf{u} \in C_t^1(\Gamma)^3, \quad (2.19)$$

$$\int_\gamma \operatorname{curl}_\Gamma \mathbf{u} \, ds = \int_{\partial\gamma} \mathbf{u} \cdot \boldsymbol{\tau} \, ds, \quad \text{for } \mathbf{u} \in C_t^1(\Gamma)^3. \quad (2.20)$$

The identity (2.19) is the fundamental Stokes result (e.g., [13]). The result in (2.20) easily follows from (2.19) and the vector-product rule also used above:

$$\int_\gamma \operatorname{curl}_\Gamma \mathbf{u} \, ds = \int_\gamma \operatorname{div}_\Gamma (\mathbf{u} \times \mathbf{n}) \, ds = \int_{\partial\gamma} (\mathbf{u} \times \mathbf{n}) \cdot \boldsymbol{\nu} \, ds = \int_{\partial\gamma} (\mathbf{n} \times \boldsymbol{\nu}) \cdot \mathbf{u} \, ds = \int_{\partial\gamma} \boldsymbol{\tau} \cdot \mathbf{u} \, ds.$$

**3. Surface Sobolev spaces.** In this section we recall and derive basic properties of surface Sobolev spaces. We will introduce  $\mathbf{H}(\operatorname{div}_\Gamma)$  and  $\mathbf{H}(\operatorname{curl}_\Gamma)$  spaces and give some properties of these spaces, which are direct analogons of well-known properties of these spaces in Euclidean  $\mathbb{R}^2$  and  $\mathbb{R}^3$ . These properties are useful in the analysis of the Helmholtz decomposition in section 4.

The Sobolev space of  $L^2(\Gamma)$  functions for which all first weak tangential derivatives are in  $L^2(\Gamma)$  can be defined by local charts as in section 4.2 in [46] and is denoted by  $H^1(\Gamma)$ . Its dual is denoted by  $H^{-1}(\Gamma)$ . Using the smoothness assumption on  $\Gamma$  it can be shown [Theorem 4.3 in [46]] that the space of smooth functions  $\mathcal{D} := C^2(\Gamma)$  is dense in  $H^1(\Gamma)$ . The space  $H^1(\Gamma)$  can also be characterized by  $H^1(\Gamma) = \overline{\mathcal{D}}^{\|\cdot\|_1}$ , where the norm  $\|\cdot\|_1$  and corresponding scalar product are given by  $(\phi, \psi)_1 := (\phi, \psi)_{L^2(\Gamma)} + (\nabla_\Gamma \phi, \nabla_\Gamma \psi)_{L^2(\Gamma)}$  and  $\|\phi\|_1^2 = (\phi, \phi)_1$ . The space of smooth vector functions on  $\Gamma$  which are tangential to the surface (hence, contained in the tangent bundle) is denoted by

$$\mathcal{D}_t^3 := \{ \mathbf{u} \in \mathcal{D}^3 \mid \mathbf{n} \cdot \mathbf{u} = 0 \text{ on } \Gamma \}.$$

We introduce the spaces of vector tangential functions

$$\begin{aligned} \mathbf{L}_t^2(\Gamma) &:= \{ \mathbf{u} \in L^2(\Gamma)^3 \mid \mathbf{n} \cdot \mathbf{u} = 0 \text{ a.e. on } \Gamma \}, \\ \mathbf{H}_t^1(\Gamma) &:= \{ \mathbf{u} \in H^1(\Gamma)^3 \mid \mathbf{n} \cdot \mathbf{u} = 0 \text{ a.e. on } \Gamma \}. \end{aligned}$$

From the density of  $\mathcal{D}$  in  $L^2(\Gamma)$  and  $H^1(\Gamma)$  it follows that  $\mathcal{D}_t^3$  is dense in both  $\mathbf{L}_t^2(\Gamma)$  and  $\mathbf{H}_t^1(\Gamma)$ . A natural norm on the latter space is  $\|\mathbf{u}\|_1^2 = \sum_{i=1}^3 \|u_i\|_1^2 = \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \sum_{i=1}^3 \|\nabla_\Gamma u_i\|_{L^2(\Gamma)}^2$ . Instead of this norm we will use another equivalent one, which is more convenient in our analysis. We derive this alternative norm. Note that

$$\sum_{i=1}^3 \|\nabla_\Gamma u_i\|_{L^2(\Gamma)}^2 \sim \|\nabla \mathbf{u} \mathbf{P}\|_{L^2(\Gamma)}^2 = \int_\Gamma \|\nabla \mathbf{u}(s) \mathbf{P}(s)\| ds,$$

where  $\|\cdot\|$  is the matrix 2-norm  $\|A\| := \rho(A^T A)^{\frac{1}{2}}$ . For  $\mathbf{u}$  that satisfies  $\mathbf{n} \cdot \mathbf{u} = 0$  on  $\Gamma$  we get (recall  $\mathbf{u} = \mathbf{u}^e$ ), with  $\mathbf{H} := \nabla_\Gamma \mathbf{n} = \nabla \mathbf{n}$  the symmetric Weingarten mapping, the relation  $\mathbf{n} \cdot \nabla \mathbf{u} = -\mathbf{u} \cdot \mathbf{H}$ . Hence,

$$\mathbf{P} \nabla \mathbf{u} \mathbf{P} = \nabla \mathbf{u} \mathbf{P} - \mathbf{n} \mathbf{n} \cdot \nabla \mathbf{u} \mathbf{P} = \nabla \mathbf{u} \mathbf{P} + \mathbf{n} \mathbf{u} \cdot \mathbf{H} \mathbf{P} = \nabla \mathbf{u} \mathbf{P} + \mathbf{n} \mathbf{u} \cdot \mathbf{H}.$$

Using  $\|\mathbf{H}\|_{L^\infty(\Gamma)} \leq c$  we obtain the norm equivalence

$$\begin{aligned} \|\mathbf{u}\|_1^2 &\sim \|\mathbf{u}\|_{\mathbf{H}^1}^2 := \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\nabla_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2 \\ &= \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \int_\Gamma \|\nabla_\Gamma \mathbf{u}\|^2 ds, \quad \mathbf{u} \in \mathbf{H}_t^1(\Gamma), \end{aligned} \tag{3.1}$$

with  $\nabla_\Gamma$  the covariant derivative  $\nabla_\Gamma \mathbf{u} = \mathbf{P} \nabla \mathbf{u} \mathbf{P}$ , cf. (2.2). In the remainder we use this norm  $\|\cdot\|_{\mathbf{H}^1}$  on  $\mathbf{H}_t^1(\Gamma)$ .

REMARK 3.1. In [21] the Poincaré inequality

$$\|\mathbf{u}\|_{L^2(\Gamma)}^2 \leq c \|\nabla_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2 \quad \text{for all } \mathbf{u} \in \mathbf{H}_t^1$$

is derived. Hence one could delete the part  $\|\cdot\|_{L^2(\Gamma)}^2$  in the norm  $\|\cdot\|_{\mathbf{H}^1}^2$ , but we will not do so.

The operators  $\nabla_\Gamma$  and  $\mathbf{curl}_\Gamma$  defined in (2.1), (2.8) are continuously extended to operators  $H^1(\Gamma) \rightarrow \mathbf{L}_t^2(\Gamma)$ . The operators  $\operatorname{div}_\Gamma, \operatorname{curl}_\Gamma$  are extended to operators  $\mathbf{L}_t^2(\Gamma) \rightarrow H^{-1}(\Gamma)$  as adjoints of  $\nabla_\Gamma$  and  $\mathbf{curl}_\Gamma$ , respectively, cf. (2.4), (2.9). In particular we have the following duality pairings:

$$\langle \operatorname{div}_\Gamma \mathbf{u}, \phi \rangle := - \int_\Gamma \mathbf{u} \cdot \nabla_\Gamma \phi \, ds \quad \text{for all } \phi \in H^1(\Gamma), \mathbf{u} \in \mathbf{L}_t^2(\Gamma), \quad (3.2)$$

$$\langle \operatorname{curl}_\Gamma \mathbf{u}, \phi \rangle := - \int_\Gamma \mathbf{u} \cdot \mathbf{curl}_\Gamma \phi \, ds \quad \text{for all } \phi \in H^1(\Gamma), \mathbf{u} \in \mathbf{L}_t^2(\Gamma). \quad (3.3)$$

Due to the density of smooth functions in  $\mathbf{H}_t^1(\Gamma)$  the identities in (2.4), (2.5), (2.9), (2.19) and (2.20) also hold with  $C^1(\Gamma)$  and  $C_t^1(\Gamma)^3$  replaced by the surface Sobolev spaces  $H^1(\Gamma)$  and  $\mathbf{H}_t^1(\Gamma)$ . The right hand side boundary integrals in (2.19) and (2.20) are then defined via the well-defined trace operator in  $H^1(\Gamma)$ .

We introduce the spaces

$$\begin{aligned} \mathbf{H}(\operatorname{div}_\Gamma) &= \{ \mathbf{u} \in \mathbf{L}_t^2(\Gamma) \mid \operatorname{div}_\Gamma \mathbf{u} \in L^2(\Gamma) \}, \quad \|\mathbf{u}\|_{\mathbf{H}(\operatorname{div}_\Gamma)}^2 = \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2, \\ \mathbf{H}(\operatorname{curl}_\Gamma) &= \{ \mathbf{u} \in \mathbf{L}_t^2(\Gamma) \mid \operatorname{curl}_\Gamma \mathbf{u} \in L^2(\Gamma) \}, \quad \|\mathbf{u}\|_{\mathbf{H}(\operatorname{curl}_\Gamma)}^2 = \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2, \\ \mathbf{X}(\Gamma) &= \mathbf{H}(\operatorname{div}_\Gamma) \cap \mathbf{H}(\operatorname{curl}_\Gamma), \quad \|\mathbf{u}\|_{\mathbf{X}}^2 = \|\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2. \end{aligned}$$

These spaces are Hilbert spaces. We will need density of smooth functions in  $\mathbf{X}(\Gamma)$ . For this we derive the following Lemma, cf. Theorems 2.4 and 2.10 in [17] for the Euclidean variant of these results. The proof given below is along the same lines as the proofs for the Euclidean case given in [17].

LEMMA 3.1. *The space  $\mathcal{D}_t^3$  is dense in  $\mathbf{H}(\operatorname{div}_\Gamma)$ ,  $\mathbf{H}(\operatorname{curl}_\Gamma)$  and  $\mathbf{X}(\Gamma)$ .*

*Proof.* The proof is based on the following elementary result:

$$\begin{aligned} \text{A subspace } M_0 \text{ of a Banach space } M \text{ is dense in } M \text{ iff} \\ \text{every element of } M' \text{ that vanishes on } M_0 \text{ also vanishes on } M. \end{aligned} \quad (3.4)$$

We first consider  $M = \mathbf{H}(\operatorname{div}_\Gamma)$ . We apply (3.4) with  $M_0 = \mathcal{D}_t^3$ . Take  $L \in \mathbf{H}(\operatorname{div}_\Gamma)'$  with  $L\mathbf{v} = 0$  for all  $\mathbf{v} \in \mathcal{D}_t^3$ . There exists a unique  $\boldsymbol{\ell} \in \mathbf{H}(\operatorname{div}_\Gamma)$  such that

$$(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} + (\operatorname{div}_\Gamma \boldsymbol{\ell}, \operatorname{div}_\Gamma \mathbf{v})_{L^2(\Gamma)} = L\mathbf{v} \quad \text{for all } \mathbf{v} \in \mathbf{H}(\operatorname{div}_\Gamma). \quad (3.5)$$

From  $L\mathbf{v} = 0$  for all  $\mathbf{v} \in \mathcal{D}_t^3$  it follows that

$$(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} = -(\operatorname{div}_\Gamma \boldsymbol{\ell}, \operatorname{div}_\Gamma \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathcal{D}_t^3. \quad (3.6)$$

Define  $\hat{D} := C^3(\Gamma)$  and note that  $\hat{D}$  is dense in  $H^2(\Gamma)$  (Theorem 4.3 in [46]). Take arbitrary  $\phi \in \hat{D}$  and  $\mathbf{v} := \nabla_\Gamma \phi \in \mathcal{D}_t^3$  in (3.6). We then get

$$(\boldsymbol{\ell}, \nabla_\Gamma \phi)_{L^2(\Gamma)} = -(\operatorname{div}_\Gamma \boldsymbol{\ell}, \Delta_\Gamma \phi)_{L^2(\Gamma)} \quad \text{for all } \phi \in \hat{D}.$$

Using  $\boldsymbol{\ell} \in \mathbf{H}(\operatorname{div}_\Gamma)$  and (3.2) it follows that

$$(\operatorname{div}_\Gamma \boldsymbol{\ell}, \phi - \Delta_\Gamma \phi)_{L^2(\Gamma)} = 0 \quad \text{for all } \phi \in \hat{D}.$$

Let  $(w_n)_{n \in \mathbb{N}} \subset H^2(\Gamma)$  be the eigensystem of the Laplace-Beltrami operator  $\Delta_\Gamma$ , with eigenvalues  $\lambda_n \geq 0$  such that  $-\Delta_\Gamma w_n = \lambda_n w_n$ . Using the density of  $\hat{D}$  in  $H^2(\Gamma)$  it follows that

$$(\operatorname{div}_\Gamma \boldsymbol{\ell}, w_n - \Delta_\Gamma w_n)_{L^2(\Gamma)} = (1 + \lambda_n)(\operatorname{div}_\Gamma \boldsymbol{\ell}, w_n)_{L^2(\Gamma)} = 0 \quad \text{for all } n \in \mathbb{N}.$$



From the density of  $(w_n)_{n \in \mathbb{N}}$  in  $L^2(\Gamma)$  it follows that  $\operatorname{div}_\Gamma \boldsymbol{\ell} = 0$  a.e. on  $\Gamma$ . Using this in (3.6) we obtain  $(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} = 0$  for all  $\mathbf{v} \in \mathcal{D}_t^3$  and due to the density of  $\mathcal{D}_t^3$  in  $\mathbf{L}_t^2(\Gamma)$  this implies  $\boldsymbol{\ell} = 0$  a.e. on  $\Gamma$ . Hence  $L$  vanishes on  $\mathbf{H}(\operatorname{div}_\Gamma)$ . This proves the density of  $\mathcal{D}_t^3$  in  $\mathbf{H}(\operatorname{div}_\Gamma)$ . With very similar arguments the density of  $\mathcal{D}_t^3$  in  $\mathbf{H}(\operatorname{curl}_\Gamma)$  can be shown. In that case we have  $\boldsymbol{\ell} \in \mathbf{H}(\operatorname{curl}_\Gamma)$  such that

$$(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} + (\operatorname{curl}_\Gamma \boldsymbol{\ell}, \operatorname{curl}_\Gamma \mathbf{v})_{L^2(\Gamma)} = L\mathbf{v} \quad \text{for all } \mathbf{v} \in \mathbf{H}(\operatorname{curl}_\Gamma), \quad (3.7)$$

and

$$(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} = -(\operatorname{curl}_\Gamma \boldsymbol{\ell}, \operatorname{curl}_\Gamma \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathcal{D}_t^3. \quad (3.8)$$

For  $\phi \in \hat{D}$  we now take  $\mathbf{v} := \mathbf{curl}_\Gamma \phi \in \mathcal{D}_t^3$ , and using (2.12) we then get

$$(\boldsymbol{\ell}, \mathbf{curl}_\Gamma \phi)_{L^2(\Gamma)} = -(\operatorname{curl}_\Gamma \boldsymbol{\ell}, \Delta_\Gamma \phi)_{L^2(\Gamma)} \quad \text{for all } \phi \in \hat{D}.$$

With the same arguments as above we can conclude  $\operatorname{curl}_\Gamma \boldsymbol{\ell} = 0$  a.e. on  $\Gamma$  and with (3.8) we get  $\boldsymbol{\ell} = 0$  a.e. on  $\Gamma$ . Hence,  $L$  vanishes on  $\mathbf{H}(\operatorname{curl}_\Gamma)$ .

The density of  $\mathcal{D}_t^3$  in the intersection  $\mathbf{X}(\Gamma)$  can also be shown by using (3.4) as follows. Take  $L \in \mathbf{X}(\Gamma)'$  with  $L\mathbf{v} = 0$  for all  $\mathbf{v} \in \mathcal{D}_t^3$ . There exists a unique  $\boldsymbol{\ell} \in \mathbf{X}(\Gamma)$  such that  $L\mathbf{v} = (\boldsymbol{\ell}, \mathbf{v})_X$  for all  $\mathbf{v} \in \mathbf{X}(\Gamma)$  and

$$(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} = -(\operatorname{div}_\Gamma \boldsymbol{\ell}, \operatorname{div}_\Gamma \mathbf{v})_{L^2(\Gamma)} - (\operatorname{curl}_\Gamma \boldsymbol{\ell}, \operatorname{curl}_\Gamma \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathcal{D}_t^3. \quad (3.9)$$

Take  $\phi \in \hat{D}$  and  $\mathbf{v} := \nabla_\Gamma \phi \in \mathcal{D}_t^3$ , hence  $\operatorname{curl}_\Gamma \mathbf{v} = 0$ . We then get  $(\boldsymbol{\ell}, \nabla_\Gamma \phi)_{L^2(\Gamma)} = -(\operatorname{div}_\Gamma \boldsymbol{\ell}, \Delta_\Gamma \phi)_{L^2(\Gamma)}$  and with the arguments used above we conclude  $\operatorname{div}_\Gamma \boldsymbol{\ell} = 0$  a.e. on  $\Gamma$ . We can also take  $\mathbf{v} := \mathbf{curl}_\Gamma \phi \in \mathcal{D}_t^3$ , hence  $\operatorname{div}_\Gamma \mathbf{v} = 0$ . We then get  $(\boldsymbol{\ell}, \mathbf{curl}_\Gamma \phi)_{L^2(\Gamma)} = -(\operatorname{curl}_\Gamma \boldsymbol{\ell}, \Delta_\Gamma \phi)_{L^2(\Gamma)}$  and from this we obtain  $\operatorname{curl}_\Gamma \boldsymbol{\ell} = 0$  a.e. on  $\Gamma$ . Using  $\operatorname{div}_\Gamma \boldsymbol{\ell} = 0$  and  $\operatorname{curl}_\Gamma \boldsymbol{\ell} = 0$  a.e. on  $\Gamma$  in (3.9) we obtain  $(\boldsymbol{\ell}, \mathbf{v})_{L^2(\Gamma)} = 0$  for all  $\mathbf{v} \in \mathcal{D}_t^3$  and with a density argument we conclude  $\boldsymbol{\ell} = 0$ , hence  $L$  vanishes on  $\mathbf{X}(\Gamma)$ .  $\square$

We now show that the spaces  $\mathbf{X}(\Gamma)$  and  $\mathbf{H}_t^1(\Gamma)$  are isomorphic.

**THEOREM 3.2.** *There are constants  $c_1, c_2$  such that*

$$\|\mathbf{u}\|_X \leq c_1 \|\mathbf{u}\|_{\mathbf{H}^1} \leq c_2 \|\mathbf{u}\|_X \quad \text{for all } \mathbf{u} \in \mathbf{X}(\Gamma) \quad (3.10)$$

holds. Hence  $\mathbf{X}(\Gamma) \simeq \mathbf{H}_t^1(\Gamma)$  holds.

*Proof.* The first estimate in (3.10) follows directly from the definition of the spaces. It suffices to prove the second inequality for the dense subspace  $\mathcal{D}_t^3$ . Take  $\mathbf{u} \in \mathcal{D}_t^3$ . Using (2.4), (2.5), (2.9) and (2.14) we get

$$\begin{aligned} & \int_\Gamma \operatorname{div}_\Gamma \mathbf{u} \operatorname{div}_\Gamma \mathbf{u} \, ds + \int_\Gamma \operatorname{curl}_\Gamma \mathbf{u} \operatorname{curl}_\Gamma \mathbf{u} \, ds \\ &= - \int_\Gamma [\nabla_\Gamma(\operatorname{div}_\Gamma \mathbf{u}) + \mathbf{curl}_\Gamma(\operatorname{curl}_\Gamma \mathbf{u})] \cdot \mathbf{u} \, ds \\ &= - \int_\Gamma [\mathbf{P} \operatorname{div}_\Gamma(\nabla_\Gamma \mathbf{u}) - K\mathbf{u}] \cdot \mathbf{u} \, ds \\ &= \int_\Gamma \operatorname{tr}((\nabla_\Gamma \mathbf{u})^T \nabla_\Gamma \mathbf{u}) + K\mathbf{u} \cdot \mathbf{u} \, ds. \end{aligned} \quad (3.11)$$

$$(3.12)$$

Using this one gets  $\|\mathbf{u}\|_{\mathbf{H}^1}^2 \leq c(\|\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2)$  and thus the second estimate in (3.10).  $\square$

REMARK 3.2. The result in Theorem 3.2 is a surface analogon of the result in Lemma 2.5 in [17]. In the latter the property  $H_0^1(\Omega)^N \simeq H_0(\operatorname{div}; \Omega) \cap H_0(\mathbf{curl}; \Omega)$  for a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^N$  is derived.

**4. Helmholtz decomposition.** In this section we derive a surface Helmholtz decomposition which states that every  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$  can be uniquely decomposed as the sum of the tangential gradient of a scalar potential, the vector surface curl of a stream function and a tangential harmonic field. We will show that if  $\Gamma$  is simply connected the harmonic field term in the decomposition must be zero. The analysis is based on elementary differential calculus and functional analysis. Concerning the latter, the main ingredients that we use are the Peetre-Tartar and Lax-Milgram lemmas.

We define the space of harmonic fields:

$$\mathcal{H} = \{ \mathbf{u} \in \mathbf{L}_t^2(\Gamma) \mid \operatorname{div}_\Gamma \mathbf{u} = 0 \quad \text{and} \quad \operatorname{curl}_\Gamma \mathbf{u} = 0 \}, \quad (4.1)$$

This is a closed subspace of  $\mathbf{L}_t^2(\Gamma)$ . Furthermore,  $\mathcal{H} \subset \mathbf{X}(\Gamma)$  holds.

LEMMA 4.1. *The space of harmonic fields has finite dimension:  $\dim(\mathcal{H}) < \infty$ .*

*Proof.* We apply a version of the Peetre-Tartar Lemma [42], which we briefly recall. Let  $E_1, E_2, E_3$  be Banach spaces,  $A : E_1 \rightarrow E_2$  linear and bounded, and  $B : E_1 \rightarrow E_3$  linear, bounded and compact. Furthermore  $\|v\|_{E_1} \simeq \|Av\|_{E_2} + \|Bv\|_{E_3}$  for all  $v \in E_1$ . Then  $\ker A$  is finite dimensional. We apply this with  $E_1 = \mathbf{X}(\Gamma)$ ,  $E_2 = L^2(\Gamma)^2$ ,  $E_3 = L^2(\Gamma)^3$ ,  $A\mathbf{u} = (\operatorname{curl}_\Gamma \mathbf{u}, \operatorname{div}_\Gamma \mathbf{u})^T$ ,  $B = \operatorname{id}$ . From the compactness of the embedding  $H^1(\Gamma) \hookrightarrow L^2(\Gamma)$  (Theorem 7.10 in [46]) it follows that  $\operatorname{id} : \mathbf{X}(\Gamma) \rightarrow L^2(\Gamma)^3$  is compact. From the definitions of the norms we get  $\|\mathbf{u}\|_X^2 = \|A\mathbf{u}\|_{L^2(\Gamma)}^2 + \|\mathbf{u}\|_{L^2(\Gamma)}^2$ . Application of the Peetre-Tartar Lemma yields the desired result.  $\square$

THEOREM 4.2 (Surface Helmholtz decomposition). *For every  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$  there exist unique  $\psi, \phi \in H_*^1(\Gamma) := \{ \phi \in H^1(\Gamma) \mid \int_\Gamma \phi \, ds = 0 \}$  and  $\boldsymbol{\xi} \in \mathcal{H}$  such that*

$$\mathbf{u} = \nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi + \boldsymbol{\xi}. \quad (4.2)$$

*The range spaces  $\nabla_\Gamma(H_*^1(\Gamma))$  and  $\mathbf{curl}_\Gamma(H_*^1(\Gamma))$  are closed in  $\mathbf{L}_t^2(\Gamma)$  and the direct sum*

$$\mathbf{L}_t^2(\Gamma) = \nabla_\Gamma(H_*^1(\Gamma)) \oplus \mathbf{curl}_\Gamma(H_*^1(\Gamma)) \oplus \mathcal{H} \quad (4.3)$$

*is  $L^2$ -orthogonal.*

*Proof.* Take  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$ . Define  $b(\psi, \xi) := \int_\Gamma \nabla_\Gamma \psi \cdot \nabla_\Gamma \xi \, ds$ . This bilinear form is continuous and elliptic on  $H_*^1(\Gamma)$ . Hence, there exists a (unique)  $\psi^* \in H_*^1(\Gamma)$  such that

$$b(\psi^*, \xi) = \int_\Gamma \mathbf{u} \cdot \nabla_\Gamma \xi \, ds \quad \text{for all } \xi \in H_*^1(\Gamma).$$

Define  $\mathbf{w} := \mathbf{u} - \nabla_\Gamma \psi^* \in \mathbf{L}_t^2(\Gamma)$ . By construction we have  $\operatorname{div}_\Gamma \mathbf{w} = 0$  in  $H^{-1}(\Gamma)$ , hence  $\mathbf{w} \in \mathbf{H}(\operatorname{div}_\Gamma)$ .

Define  $\tilde{b}(\phi, \xi) := \int_\Gamma \mathbf{curl}_\Gamma \phi \cdot \mathbf{curl}_\Gamma \xi \, ds$ . Using (2.12) it follows that  $\tilde{b}(\phi, \xi) = b(\phi, \xi)$  for all  $\phi, \xi \in H^1(\Gamma)$  and thus also  $\tilde{b}(\cdot, \cdot)$  is continuous and elliptic on  $H_*^1(\Gamma)$ . There exists a (unique)  $\phi^* \in H_*^1(\Gamma)$  such that

$$\tilde{b}(\phi^*, \xi) = \int_\Gamma \mathbf{w} \cdot \mathbf{curl}_\Gamma \xi \, ds \quad \text{for all } \xi \in H_*^1(\Gamma).$$

By construction we have  $\operatorname{curl}_\Gamma(\mathbf{w} - \mathbf{curl}_\Gamma\phi^*) = 0$  in  $H^{-1}(\Gamma)$ , hence  $\mathbf{w} - \mathbf{curl}_\Gamma\phi^* \in \mathbf{H}(\operatorname{curl}_\Gamma)$ . Note that

$$\langle \operatorname{div}_\Gamma \mathbf{curl}_\Gamma\phi, \xi \rangle = - \int_\Gamma \mathbf{curl}_\Gamma\phi \cdot \nabla_\Gamma\xi \, ds = 0 \quad \text{for all } \phi, \xi \in H^1(\Gamma). \quad (4.4)$$

Define  $\xi := \mathbf{u} - \nabla_\Gamma\psi^* - \mathbf{curl}_\Gamma\phi^* = \mathbf{w} - \mathbf{curl}_\Gamma\phi^*$ . Using (4.4) we obtain  $\operatorname{div}_\Gamma\xi = \operatorname{div}_\Gamma\mathbf{w} = 0$  in  $H^{-1}(\Gamma)$ . We also have  $\operatorname{curl}_\Gamma\xi = 0$  in  $H^{-1}(\Gamma)$ . Thus  $\xi \in \mathcal{H}$ . Hence we have a representation of  $\mathbf{u}$  as in (4.2). From the Poincare inequality  $\|\psi\|_1 \leq c\|\nabla_\Gamma\psi\|_{L^2(\Gamma)}$  for all  $\psi \in H_*^1(\Gamma)$  it follows that the range space  $\nabla_\Gamma(H_*^1(\Gamma))$  is closed in  $\mathbf{L}_t^2(\Gamma)$  and that  $\nabla_\Gamma : H_*^1(\Gamma) \rightarrow \mathbf{L}_t^2(\Gamma)$  is injective. From this and  $\|\mathbf{curl}_\Gamma\phi\|_{L^2(\Gamma)} = \|\nabla_\Gamma\phi\|_{L^2(\Gamma)}$  it follows that the range space  $\mathbf{curl}_\Gamma(H_*^1(\Gamma))$  is closed in  $\mathbf{L}_t^2(\Gamma)$  and that  $\mathbf{curl}_\Gamma : H_*^1(\Gamma) \rightarrow \mathbf{L}_t^2(\Gamma)$  is injective. The orthogonality of the decomposition in (4.3) easily follows from (4.4). The uniqueness of  $\psi$ ,  $\phi$  and  $\xi$  in (4.2) follows from the orthogonality property and the injectivity of  $\nabla_\Gamma$  and  $\mathbf{curl}_\Gamma$ .  $\square$

For the formulation of the Stokes problem in rotation formulation, treated in section 5, it is essential that there are no nontrivial harmonic fields, i.e.,  $\dim(\mathcal{H}) = 0$ . This result holds provided the surface  $\Gamma$  is *simply connected* and can be derived using elementary calculus. This derivation is given in Lemma 4.3 below. If  $\Gamma$  is *not* simply connected but has a genus  $> 1$ , then  $\dim(\mathcal{H}) > 0$  and  $\dim(\mathcal{H})$  can be directly related to the genus, cf. Remark 4.2.

LEMMA 4.3. *Assume that  $\Gamma$  is simply connected. Then  $\dim(\mathcal{H}) = 0$  holds.*

*Proof.* Take  $\mathbf{u} \in \mathcal{H}$ . Hence  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$ ,  $\operatorname{div}_\Gamma\mathbf{u} = 0$ ,  $\operatorname{curl}_\Gamma\mathbf{u} = 0$ . This implies  $\mathbf{u} \in \mathbf{X}(\Gamma)$  and due to (3.10) we get  $\mathbf{u} \in \mathbf{H}_t^1(\Gamma)$ . From elliptic regularity theory as in e.g. [28] it follows that, provided  $\Gamma$  is sufficiently smooth, we have  $\mathbf{u} \in C(\Gamma)^3$ . To make this more precise we note the following. We have  $\mathbf{u} \in \mathcal{H}$  iff  $D(\mathbf{u}) := (\operatorname{div}_\Gamma\mathbf{u}, \operatorname{div}_\Gamma\mathbf{u})_{L^2(\Gamma)} + (\operatorname{curl}_\Gamma\mathbf{u}, \operatorname{curl}_\Gamma\mathbf{u})_{L^2(\Gamma)} = 0$ . This Dirichlet integral  $D(\mathbf{u})$  corresponds to the Hodge Laplacian, cf. (4.10) below, which is an elliptic operator. This ellipticity can also be concluded from the relation (3.12). From elliptic regularity theory, e.g., the result (vi) on page 296 in [28], it follows that if  $\Gamma$  has Hölder smoothness  $C_\mu^k$  ( $k \in \mathbb{N}$ ,  $0 < \mu \leq 1$ ) then the harmonic fields  $\mathbf{u} \in \mathcal{H}$  have Hölder smoothness  $\mathbf{u} \in C_\mu^{k-1}$  (componentwise). Using assumption 2.1 we conclude that  $\mathbf{u} \in C(\Gamma)^3$ .

For a (piecewise) regular parametrized differentiable curve (cf. [11])  $\alpha : [a, b] \subset \mathbb{R} \rightarrow \Gamma$  we denote the line integral of a function  $f : \operatorname{im}(\alpha) \rightarrow \mathbb{R}$  by

$$\int_\alpha f \, ds := \int_a^b f(\alpha(t)) \|\alpha'(t)\| \, dt.$$

A parametrized curve  $g(t)$  on  $\Gamma$  is called a geodesic if the covariant derivative of the vector field  $g'(t)$  along  $\operatorname{im}(g)$  equals zero. The latter property is equivalent to the condition that  $g''(t)$  is orthogonal to  $\Gamma$ . We take an arbitrary fixed point  $x_0$  on  $\Gamma$ . From the Hopf-Rinow theorem (cf. [11]) it follows that for all  $x \in \Gamma$ ,  $x \neq x_0$ , there exists a minimal (i.e., length minimizing) geodesic, which is denoted by  $g_x(t)$ . This  $g_x$  may be non-unique. For the given  $\mathbf{u} \in \mathcal{H}$  (note that  $\mathbf{u} \in C(\Gamma)^3$ ) we define

$$\psi(x) := \int_{g_x} \mathbf{u} \cdot \frac{g'_x}{\|g'_x\|} \, ds \quad \text{for } x \in \Gamma, x \neq x_0, \quad \psi(x_0) := 0. \quad (4.5)$$

We now show that this definition of  $\psi$  does not depend on the particular choice of  $g_x$ . A generic situation with two different minimal geodesics  $g_x$  and  $\tilde{g}_x$  is sketched in Fig. 4.1.




Fig. 4.1: Multiple minimal geodesics (left). Tangential derivative (right).

Due to the essential assumption that  $\Gamma$  is simply connected, the domain  $\gamma$  enclosed by the curves  $g_A$  and  $\tilde{g}_A$  is contained in  $\Gamma$ . Using the same notation  $\boldsymbol{\tau} = \mathbf{n} \times \boldsymbol{\nu}$  for the (oriented) tangential vector on  $\partial\gamma$  as in (2.20) we have  $\frac{g'_A}{\|g'_A\|} = \pm\boldsymbol{\tau}$ , where the sign depends on the orientation of  $\Gamma$ . Without loss of generality we can assume the “+” sign. We then also have  $\frac{\tilde{g}'_A}{\|\tilde{g}'_A\|} = -\boldsymbol{\tau}$  on  $\text{im}(\tilde{g}_A)$ . Using this and

$$\int_{\partial\gamma} \mathbf{u} \cdot \boldsymbol{\tau} ds = \int_{\gamma} \text{curl}_{\Gamma} \mathbf{u} ds = 0$$

we get

$$\int_{g_A} \mathbf{u} \cdot \frac{g'_A}{\|g'_A\|} ds = \int_{\tilde{g}_A} \mathbf{u} \cdot \frac{\tilde{g}'_A}{\|\tilde{g}'_A\|} ds.$$

The same argument can be applied for the minimal geodesics connecting  $A$  and  $x$ , cf. Fig. 4.1. Hence, the definition of  $\psi$  in (4.5) does not depend on the choice of the minimal geodesic  $g_x$ .

We now consider the tangential derivative of  $\psi$  at  $x \in \Gamma$ . We assume  $x \neq x_0$ . Let  $g_x$  be a minimal geodesic connecting  $x_0$  and  $x$ , and  $t_1$  the parameter value such that  $g_x(t_1) = x$ . Define  $\mathbf{w} := g'_x(t_1)$ . Take  $\mathbf{a} \in T_x\Gamma$  (tangential plane at  $x$ ),  $\mathbf{a} \neq 0$ . We assume that  $\mathbf{a}$  and  $\mathbf{w}$  are linearly independent, cf. Fig. 4.1. Let  $\alpha$  be the unique geodesic with  $\alpha(0) = x$ ,  $\alpha'(0) = \mathbf{a}$ , cf., e.g., Chapter 7 in [45]. For  $\epsilon > 0$  sufficiently small the geodesics  $g_x$  and  $g_{\alpha(\epsilon)}$  do not intersect. Using (2.20),  $\text{curl}_{\Gamma} \mathbf{u} = 0$  and  $\tilde{\gamma} \subset \Gamma$  (cf. Fig. 4.1 for notation) we have  $\int_{\partial\tilde{\gamma}} \mathbf{u} \cdot \boldsymbol{\tau} ds = 0$  and thus we get

$$\begin{aligned} \nabla_{\Gamma} \psi(x) \cdot \mathbf{a} &= \lim_{\epsilon \downarrow 0} \frac{\psi(\alpha(\epsilon)) - \psi(x)}{\epsilon} \\ &= \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \left[ \int_{g(\alpha(\epsilon))} \mathbf{u} \cdot \frac{g'_{\alpha(\epsilon)}}{\|g'_{\alpha(\epsilon)}\|} ds - \int_{g_x} \mathbf{u} \cdot \frac{g'_x}{\|g'_x\|} ds \right] \\ &= \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \int_{\alpha(\epsilon)} \mathbf{u} \cdot \frac{\alpha'}{\|\alpha'\|} ds = \lim_{\epsilon \downarrow 0} \frac{1}{\epsilon} \int_0^{\epsilon} \mathbf{u}(\alpha(t)) \cdot \alpha'(t) dt = \mathbf{u}(x) \cdot \mathbf{a}. \end{aligned}$$

Hence  $\nabla_{\Gamma} \psi(x) \cdot \mathbf{a} = \mathbf{u}(x) \cdot \mathbf{a}$  if  $\mathbf{a}$  and  $\mathbf{w}$  are not linearly dependent. With very similar arguments one can show that the same identity holds if  $\mathbf{a}$  and  $\mathbf{w}$  are linearly dependent. We conclude that  $\nabla_{\Gamma} \psi(x) = \mathbf{u}(x)$  for all  $x \neq x_0$ . One may check that the arguments above also apply for  $x = x_0$  (i.e.,  $\psi(x) = 0$ ). Hence,  $\nabla_{\Gamma} \psi = \mathbf{u}$  on  $\Gamma$ .

From  $\operatorname{div}_\Gamma \mathbf{u} = 0$  we then obtain  $\Delta_\Gamma \psi = 0$  on  $\Gamma$ . Hence,  $\psi$  must be a constant on  $\Gamma$ . Consequently  $\nabla_\Gamma \psi = \mathbf{u} = 0$  on  $\Gamma$ , which completes the proof.  $\square$

We finally formulate two corollaries.

**COROLLARY 4.4.** *Let  $\Gamma$  be simply connected. For the operators  $\operatorname{curl}_\Gamma, \operatorname{div}_\Gamma : \mathbf{L}_t^2(\Gamma) \rightarrow H^{-1}(\Gamma)$  and  $\mathbf{curl}_\Gamma, \nabla_\Gamma : H^1(\Gamma) \rightarrow \mathbf{L}_t^2(\Gamma)$  the following holds:*

$$\ker(\operatorname{div}_\Gamma) = \operatorname{im}(\mathbf{curl}_\Gamma), \quad (4.6)$$

$$\ker(\operatorname{curl}_\Gamma) = \operatorname{im}(\nabla_\Gamma). \quad (4.7)$$

*Proof.* We consider (4.6). Take  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$  and its Helmholtz decomposition  $\mathbf{u} = \nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi$ , with unique  $\psi, \phi \in H_*^1(\Gamma)$ . Now note (cf. (4.4))

$$\begin{aligned} \operatorname{div}_\Gamma \mathbf{u} = 0 &\Leftrightarrow (\mathbf{u}, \nabla_\Gamma \xi)_{L^2(\Gamma)} = 0 \quad \text{for all } \xi \in H^1(\Gamma) \\ &\Leftrightarrow (\nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi, \nabla_\Gamma \xi)_{L^2(\Gamma)} = 0 \quad \text{for all } \xi \in H^1(\Gamma) \\ &\Leftrightarrow (\nabla_\Gamma \psi, \nabla_\Gamma \xi)_{L^2(\Gamma)} = 0 \quad \text{for all } \xi \in H^1(\Gamma) \\ &\Leftrightarrow \psi = 0 \\ &\Leftrightarrow \mathbf{u} \in \operatorname{im}(\mathbf{curl}_\Gamma). \end{aligned}$$

The result in (4.7) follows with similar arguments or by noting that  $\operatorname{curl}_\Gamma, \nabla_\Gamma$  are (minus) the adjoints of  $\mathbf{curl}_\Gamma$  and  $\operatorname{div}_\Gamma$ , respectively.  $\square$

**COROLLARY 4.5** (Friedrichs inequality). *Assume that  $\Gamma$  is simply connected. There exists a constant  $c$  such that*

$$\|\mathbf{u}\|_{\mathbf{H}^1}^2 \leq c(\|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2 + \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}^2) \quad \text{for all } \mathbf{u} \in \mathbf{H}_t^1(\Gamma).$$

*Proof.* We use the Helmholtz decomposition as in (4.2) with  $\boldsymbol{\xi} = 0$ , i.e.  $\mathbf{u} = \nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi$  and  $\|\mathbf{u}\|_{L^2(\Gamma)}^2 = \|\nabla_\Gamma \psi\|_{L^2(\Gamma)}^2 + \|\mathbf{curl}_\Gamma \phi\|_{L^2(\Gamma)}^2$ . Using this, the result (4.7) and the Friedrichs inequality in  $H_*^1(\Gamma)$  we get

$$\begin{aligned} \|\nabla_\Gamma \psi\|_{L^2(\Gamma)}^2 &= \int_\Gamma \mathbf{u} \cdot \nabla_\Gamma \psi \, ds - \int_\Gamma \mathbf{curl}_\Gamma \phi \cdot \nabla_\Gamma \psi \, ds = - \int_\Gamma \operatorname{div}_\Gamma \mathbf{u} \psi \, ds \\ &\leq \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)} \|\psi\|_{L^2(\Gamma)} \leq c \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)} \|\nabla_\Gamma \psi\|_{L^2(\Gamma)}. \end{aligned}$$

Hence,  $\|\nabla_\Gamma \psi\|_{L^2(\Gamma)} \leq c \|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}$  holds. With similar arguments one obtains  $\|\mathbf{curl}_\Gamma \phi\|_{L^2(\Gamma)} \leq c \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}$ . Thus we get

$$\|\mathbf{u}\|_{L^2(\Gamma)} \leq c(\|\operatorname{div}_\Gamma \mathbf{u}\|_{L^2(\Gamma)} + \|\operatorname{curl}_\Gamma \mathbf{u}\|_{L^2(\Gamma)}),$$

and combining this with the upper bound in (3.10) yields the result.  $\square$

**REMARK 4.1.** We relate some of the results derived in this section to well-known fundamental results for the Euclidean case, i.e., for a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^N$ . If  $\Gamma$  is simply connected then the Helmholtz decomposition (4.2) (with  $\boldsymbol{\xi} = 0$ ) implies the following: a function  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$  satisfies  $\operatorname{curl}_\Gamma \mathbf{u} = 0$  on  $\Gamma$  iff there exists a unique  $\psi \in H_*^1(\Gamma)$  such that  $\mathbf{u} = \nabla_\Gamma \psi$ . This is the analogon of the result in [17] Theorem 2.9. From Corollary 4.4 we obtain:

$$\mathbf{L}_t^2(\Gamma) = \ker(\operatorname{div}_\Gamma) \oplus \ker(\operatorname{div}_\Gamma)^\perp = \ker(\operatorname{div}_\Gamma) \oplus \operatorname{im}(\nabla_\Gamma) = \ker(\operatorname{div}_\Gamma) \oplus \ker(\operatorname{curl}_\Gamma),$$

which is the analogon of  $L^2(\Omega)^N = H \oplus H^\perp$  (page 29 in [17]) and Corollary 2.9 in [17]. The Euclidean variant of the Friedrichs inequality in Corollary 4.5 is discussed in Remark 3.5 in [17]. In Theorem 3.1 [17] the following fundamental result is derived (where  $\Gamma_i$ ,  $0 \leq i \leq p$ , are the boundary components of the possibly multiply connected domain  $\Omega \subset \mathbb{R}^2$ ): a function  $\mathbf{v} \in L^2(\Omega)^2$  satisfies [  $\operatorname{div} \mathbf{v} = 0$  and  $\langle \mathbf{v} \cdot \mathbf{n} \rangle_{\Gamma_i} = 0$ ,  $0 \leq i \leq p$  ] iff [ there exists a stream function  $\phi \in H^1(\Omega)$  such that  $\mathbf{v} = \mathbf{curl} \phi$  ]. An analogous result in our setting follows from the Helmholtz decomposition in Theorem 4.2: [  $\operatorname{div}_\Gamma \mathbf{u} = 0$  and  $\mathbf{u} \perp \mathcal{H}$  ] iff [ there exists a stream function  $\phi \in H_*^1(\Gamma)$  such that  $\mathbf{u} = \mathbf{curl}_\Gamma \phi$  ]. Note that in case of a simply connected domain  $\Omega$  (i.e.,  $p = 0$ ) and a simply connected  $\Gamma$  the condition  $\langle \mathbf{v} \cdot \mathbf{n} \rangle_{\Gamma_0} = 0$  follows from  $\operatorname{div} \mathbf{v} = 0$  (and thus can be deleted) and  $\mathbf{u} \perp \mathcal{H}$  is automatically satisfied due to  $\mathcal{H} = \{0\}$ . Finally we note that different versions of the Helmholtz decomposition (in Euclidean space) exist. One version is given in Theorem 3.2 in [17]. This version and a comparison with various variants is given in [10]. The surface Helmholtz decomposition in Theorem 4.2 is the analogon of the following Euclidean version given in Theorem 13 in [10]:  $L^2(\Omega)^2 = X_0 \oplus W_0 \oplus R$ , with  $X_0 = \{ \nabla \psi \mid \psi \in H_0^1(\Omega) \}$ ,  $W_0 = \{ \mathbf{curl} \phi \mid \phi \in H_0^1(\Omega) \}$  and  $R = \{ \mathbf{v} \in L^2(\Omega)^2 \mid \operatorname{div} \mathbf{v} = 0 \text{ and } \operatorname{curl} \mathbf{v} = 0 \}$ .

**4.1. Relation to Hodge decomposition.** As is known from the literature, the Helmholtz decomposition can be seen as a special case of the much more general Hodge decomposition, which is derived in the framework of differential forms. In this section we derive and discuss some relevant relations between the surface differential operators and the Helmholtz decomposition introduced above and analogous notions and results known in the field of differential forms. The discussion on this topic is not essential for the results derived in Sections 5-6.

We make use of the exposition given in the Appendix of [9]. The presentation in this reference is very useful for us, because it emphasizes relevant relations between operators from differential geometry and the surface differential operators introduced above. We only outline a few results that are relevant for the discussion here. In particular we give results for the case of a 2-dimensional surface without boundary embedded in  $\mathbb{R}^3$ . We use the notation from [9] (Appendix, Sect. 6.2). For precise definitions and more detailed explanations we refer to [9]. The tangent and cotangent bundles are denoted by  $T\Gamma = \cup_{x \in \Gamma} T_x \Gamma$  and  $T^*\Gamma = \cup_{x \in \Gamma} T_x^* \Gamma$ . In the domain of a local coordinate system  $(x^1, x^2)$  (corresponding to a local parametrization) basis vectors of the tangent space  $T_x \Gamma$  at  $x \in \Gamma$  are denoted by  $(\partial x^1)_x, (\partial x^2)_x$  and the associated dual basis of  $T_x^* \Gamma$  is denoted by  $(dx^1)_x, (dx^2)_x$ . The metric is defined by the Euclidean scalar product in  $\mathbb{R}^3$ , i.e., the first fundamental form  $g : \Gamma \rightarrow T^*\Gamma \times T^*\Gamma$  is  $g_x(\mathbf{v}, \mathbf{w}) = \langle \mathbf{v}, \mathbf{w} \rangle$ , for  $x \in \Gamma$ ,  $\mathbf{v}, \mathbf{w} \in T_x \Gamma$  and  $\langle \cdot, \cdot \rangle$  the Euclidean scalar product in  $\mathbb{R}^3$ . The operator representation of the bilinear form  $g_x(\cdot, \cdot)$  is denoted by  $G_x$ , i.e.,  $G_x : T_x \Gamma \rightarrow T_x^* \Gamma$  is defined by  $G_x(\mathbf{v})(\mathbf{w}) = g_x(\mathbf{v}, \mathbf{w}) = \langle \mathbf{v}, \mathbf{w} \rangle$ . For the 1-form  $G_x(\mathbf{v}) \in T_x^* \Gamma$  the notation  $\omega_{\mathbf{v}}$  is used (note that the dependence on  $x$  is dropped in the notation). The area 2-form associated to  $g$  is given by  $v^g := \pm dx^1 \wedge dx^2 \mid \det g \mid^{\frac{1}{2}}$  (sign depending on the orientation of  $\Gamma$ ). Functions on  $\Gamma$  are called 0-forms. On the spaces of 0- and 1-forms we introduce the scalar products

$$(f, h) = \int_{\Gamma} f h \, d\Gamma \quad f, h \in L^2(\Gamma), \quad (\omega, \eta) = \int_{\Gamma} \omega_x (G_x^{-1} \eta_x) \, d\Gamma \quad \omega, \eta \in T^*\Gamma,$$

where  $d\Gamma$  is the surface measure induced by  $g$ . An analogous scalar product is used on the space of 2-forms. The space  $L_r^2(\Gamma)$  ( $r = 0, 1, 2$ ) is the closure of the space of smooth differential  $r$ -forms (note that  $L_0^2(\Gamma) = L^2(\Gamma)$ ). The Hodge transformation,

denoted by  $*$ , maps  $r$ -forms to  $(2-r)$ -forms ( $r = 0, 1, 2$ ) and is an isometry  $*$  :  $L_r^2(\Gamma) \rightarrow L_{2-r}^2(\Gamma)$ . Note that  $*v^g = 1$ ,  $*1 = v^g$ . The *exterior derivative*, which maps an  $r$ -form to an  $(r+1)$ -form ( $r = 0, 1$ ) is denoted by  $d$ . For a smooth function  $f$  (in the local coordinate system  $(x^1, x^2)$ ) we have  $df = \sum_{i=1}^2 \frac{\partial f}{\partial x^i} dx^i$  and for a 1-form  $\omega = \sum_{i=1}^2 \omega_i dx^i$ , with coefficient functions  $\omega_i$ , we have  $d\omega = \sum_{i,j=1}^2 \frac{\partial \omega_i}{\partial x^j} dx^i \wedge dx^j$ . For an  $r$ -form  $\omega$  ( $r = 1, 2$ ) the *codifferential*  $\delta\omega$  is an  $(r-1)$ -form defined by  $\delta\omega = -*dast\omega$ . The operator  $\delta$  is the adjoint of  $d$ :

$$(df, \alpha) = (f, \delta\alpha) \quad \text{for 0-forms } f, \text{ 1-forms } \alpha. \quad (4.8)$$

There are basic relations between  $d$ ,  $\delta$  (applied to differential forms) and the differential operators defined in section 2, which we now discuss. For a smooth function  $f$  we define  $\nabla_\Gamma f := G^{-1}df$ ; one easily checks that  $\nabla_\Gamma f$  is the same as the tangential gradient defined in (2.1). Further canonical definitions are (with a tangential vector field  $\mathbf{u} \in T\Gamma$ ):

$$\operatorname{div}_\Gamma \mathbf{u} := -\delta\omega_{\mathbf{u}}, \quad \operatorname{curl}_\Gamma \mathbf{u} := *d\omega_{\mathbf{u}}, \quad \mathbf{curl}_\Gamma f := -G^{-1}\delta(fv^g). \quad (4.9)$$

From this one can derive the relations (cf. [9]):

$$d\omega_{\mathbf{u}} = (\operatorname{curl}_\Gamma \mathbf{u})v^g, \quad \delta(fv^g) = -\omega_{\mathbf{curl}_\Gamma f}.$$

For the  $\operatorname{div}_\Gamma$  operator defined in (4.9) we obtain, using (4.8), for arbitrary (smooth) functions  $f$ :

$$\begin{aligned} \int_\Gamma \operatorname{div}_\Gamma \mathbf{u} f \, d\Gamma &= -(\delta\omega_{\mathbf{u}}, f) = -(\omega_{\mathbf{u}}, df) = -\int_\Gamma \omega_{\mathbf{u}}(G^{-1}df) \, d\Gamma \\ &= -\int_\Gamma \omega_{\mathbf{u}}(\nabla_\Gamma f) \, d\Gamma = -\int_\Gamma \langle \mathbf{u}, \nabla_\Gamma f \rangle \, d\Gamma, \end{aligned}$$

and comparing this with (2.4) it follows that this operator  $\operatorname{div}_\Gamma$  is the same as the one defined in (2.3) (namely minus the adjoint of  $\nabla_\Gamma$ ). With similar basic arguments (cf. [9]) one can derive for the  $\operatorname{curl}_\Gamma$  and  $\mathbf{curl}_\Gamma$  operators defined in (4.9) the relations

$$\operatorname{curl}_\Gamma \mathbf{u} = \operatorname{div}_\Gamma(\mathbf{u} \times \mathbf{n}), \quad \mathbf{curl}_\Gamma f = \mathbf{n} \times \nabla_\Gamma f,$$

hence these operators are the same as the ones defined in (2.7), (2.8). The *Hodge Laplacian* is defined by  $\Delta^H := -(d\delta + \delta d)$  and maps  $r$ -forms to  $r$ -forms ( $r = 0, 1, 2$ ). For  $r = 0$  we have

$$\Delta^H f = -\delta df = -\delta(G\nabla_\Gamma f) = -\delta(\omega_{\nabla_\Gamma f}) = \operatorname{div}_\Gamma \nabla_\Gamma f = \Delta_\Gamma f.$$

Application to a 1-form yields:

$$\begin{aligned} \Delta^H \omega_{\mathbf{u}} &= -(d\delta + \delta d)\omega_{\mathbf{u}} = d(\operatorname{div}_\Gamma \mathbf{u}) - \delta(\operatorname{curl}_\Gamma \mathbf{u} v^g) \\ &= G(\nabla_\Gamma \operatorname{div}_\Gamma \mathbf{u}) + \omega_{\mathbf{curl}_\Gamma \operatorname{curl}_\Gamma \mathbf{u}} = G((\nabla_\Gamma \operatorname{div}_\Gamma + \mathbf{curl}_\Gamma \operatorname{curl}_\Gamma) \mathbf{u}). \end{aligned}$$

Hence, the corresponding Hodge Laplacian for vector fields is given by

$$\tilde{\Delta}^H := G^{-1}\Delta^H G = \nabla_\Gamma \operatorname{div}_\Gamma + \mathbf{curl}_\Gamma \operatorname{curl}_\Gamma. \quad (4.10)$$

From (2.14) we obtain the identity

$$\tilde{\Delta}^H \mathbf{u} = \mathbf{P} \operatorname{div}_\Gamma(\nabla_\Gamma \mathbf{u}) - K\mathbf{u} = \Delta^B \mathbf{u} - K\mathbf{u}, \quad (4.11)$$

where  $\Delta^B := \mathbf{P} \operatorname{div}_\Gamma \nabla_\Gamma$  is the so-called Bochner Laplacian. The relation (4.11) corresponds to the so-called Weitzenböck identity in differential geometry, which relates the Bochner Laplacian to the Hodge Laplacian. Note that in the definition of the Bochner Laplacian the divergence operator  $\operatorname{div}_\Gamma$  applied to a matrix valued function as defined in (2.3) is used, which has no natural analogon in the setting of differential forms.

We summarize the Hodge decomposition for the special case of  $r$ -forms on a the two-dimensional surface  $\Gamma$  and then relate it to the Helmholtz decomposition derived in Theorem 4.2. Define

$$\begin{aligned} H(d, \Gamma) &:= \{ f \in L_0^2(\Gamma) \mid df \in L_1^2(\Gamma) \} \\ H(\delta, \Gamma) &:= \{ v \in L_2^2(\Gamma) \mid \delta v \in L_1^2(\Gamma) \} \\ H_1(\Gamma) &:= \{ \omega \in H_1^1(\Gamma) \mid d\omega = 0 \text{ and } \delta\omega = 0 \} \end{aligned} \quad (4.12)$$

(where  $H_1^1(\Gamma)$  is a Sobolev space of 1-forms). The space  $H_1(\Gamma)$  in (4.12) is called the *space of 1-harmonics*. The Hodge decomposition is described in the following theorem (theorems 12, 13 in Appendix of [9]).

**THEOREM 4.6.** *The spaces  $\operatorname{im} d := dH(d, \Gamma)$  and  $\operatorname{im} \delta := \delta H(\delta, \Gamma)$  are closed subspaces of  $L_1^2(\Gamma)$ ,  $\dim(H_1(\Gamma)) < \infty$  holds, and there is an  $L^2$ -orthogonal decomposition*

$$L_1^2(\Gamma) = \operatorname{im} d \oplus \operatorname{im} \delta \oplus H_1(\Gamma). \quad (4.13)$$

For  $\omega \in L_1^2(\Gamma)$  consider a decomposition

$$\omega = df + \delta v + \alpha, \text{ with } f \in H(d, \Gamma), v \in H(\delta, \Gamma), \alpha \in H_1(\Gamma). \quad (4.14)$$

Then  $\alpha$  is uniquely determined, but  $f$  and  $v$  are in general not unique. For  $f$  and  $v$  one can take  $f = \delta\omega_0$ ,  $v = d\omega_0$ , where  $\omega_0$  is the unique solution of the variational formulation of the elliptic problem  $-\Delta^H \omega_0 = \omega - \alpha$  in the Sobolev space  $V_1 := \{ \omega \in H_1^1(\Gamma) \mid \omega \text{ is } L^2\text{-orthogonal to } H_1(\Gamma) \}$ .

The decomposition in (4.13) can be directly related to the Helmholtz decomposition in (4.3). Take a decomposition of  $\mathbf{u} \in \mathbf{L}_t^2(\Gamma)$  as in (4.2) and note that

$$\begin{aligned} \mathbf{u} &= \nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi + \boldsymbol{\xi} \quad \text{iff } \omega_{\mathbf{u}} = G \nabla_\Gamma \psi + G \mathbf{curl}_\Gamma \phi + G \boldsymbol{\xi} \\ &\text{iff } \omega_{\mathbf{u}} = d\psi - \delta(\phi v^g) + \omega_{\boldsymbol{\xi}}, \end{aligned}$$

and  $[\operatorname{div}_\Gamma \boldsymbol{\xi} = 0 \text{ and } \operatorname{curl}_\Gamma \boldsymbol{\xi} = 0] \text{ iff } [\delta\omega_{\boldsymbol{\xi}} = 0 \text{ and } d\omega_{\boldsymbol{\xi}} = 0]$ . This shows the correspondence of the decompositions. Note that in (4.2) we have uniqueness of the functions  $\psi$ ,  $\phi$ , which in general does not hold in (4.14).

In the setting of differential forms an important result concerning  $\dim(H_1(\Gamma))$  can be derived. For this we recall the definition of the first de Rham cohomology group. A 1-form  $\omega$  is called closed if  $d\omega = 0$  and it is called exact if  $\omega \in \operatorname{im} d$ . The first de Rham cohomology group  $H_{dR}^1(\Gamma)$  consists of the set of (smooth) closed 1-forms modulo the exact ones. From the Hodge decomposition it easily follows that  $H_{dR}^1(\Gamma) \cong H_1(\Gamma)$ . The dimension of the first de Rham cohomology group is called the *first Betti number*  $b_1(\Gamma) := \dim(H_{dR}^1(\Gamma))$ . Extensive analysis and results for the de Rham cohomology are available, cf. e.g. [6, 25]. For example,  $H_{dR}^1(\Gamma)$  and thus also  $b_1(\Gamma)$  are homotopy invariant.

**REMARK 4.2.** The Betti number depends only on the topology of the surface. For arbitrary connected closed orientable surfaces  $\Gamma$  the value of the corresponding



first Betti number  $b_1(\Gamma)$  is known. An interesting relation is (for two-dimensional connected closed orientable surfaces)  $b_1(\Gamma) = 2 - \chi_\Gamma = 2g$ , where  $\chi_\Gamma$  is the Euler characteristic and  $g$  the genus of  $\Gamma$ . The classification theorem of such surfaces, cf. e.g. [15], yields that  $\Gamma$  is homeomorphic to either a sphere or an  $n$ -torus (connected sum of  $n$  tori, having  $n$  holes). If  $\Gamma$  is simply connected, e.g. a sphere, then  $b_1(\Gamma) = 0$  holds (which also follows from Lemma 4.3). If  $\Gamma$  is the  $n$ -torus then  $b_1(\Gamma) = 2n$ .

**5. Surface Stokes problem in stream function formulation.** In this section we consider a stationary surface Stokes problem. This problem will be reformulated in an equivalent stream function formulation. Well-posedness of the latter formulation will be discussed. As already noted above, cf. (4.11), different surface vector Laplacians are used in the literature. For the Stokes problem studied in this paper we use a Laplacian that is motivated by the modeling of surface fluids, studied in e.g., [19, 5, 23, 22, 27]. In these models the following surface rate-of-strain tensor [19] is used:

$$E_s(\mathbf{u}) := \frac{1}{2}\mathbf{P}(\nabla\mathbf{u} + \nabla\mathbf{u}^T)\mathbf{P} = \frac{1}{2}(\nabla_\Gamma\mathbf{u} + \nabla_\Gamma\mathbf{u}^T). \quad (5.1)$$

For a given force vector  $\mathbf{f} \in L^2(\Gamma)^3$ , with  $\mathbf{f} \cdot \mathbf{n} = 0$  we consider the surface Stokes problem: Find the fluid velocity tangential vector field  $\mathbf{u} : \Gamma \rightarrow \mathbb{R}^3$ , with  $\mathbf{u} \cdot \mathbf{n} = 0$ , and the surface fluid pressure  $p$  such that

$$-\mathbf{P} \operatorname{div}_\Gamma(E_s(\mathbf{u})) + \nabla_\Gamma p = \mathbf{f} \quad \text{on } \Gamma, \quad (5.2)$$

$$\operatorname{div}_\Gamma \mathbf{u} = 0 \quad \text{on } \Gamma. \quad (5.3)$$

From problem (5.2)-(5.3) one readily observes the following: the pressure field is defined up a hydrostatic mode and all tangentially rigid surface fluid motions, i.e. satisfying  $E_s(\mathbf{u}) = 0$ , are in the kernel of the differential operators on the left hand side of eq. (5.2). Integration by parts implies a consistency condition for the right hand side of eq. (5.2):

$$\int_\Gamma \mathbf{f} \cdot \mathbf{v} \, ds = 0 \quad \text{for all smooth tangential vector fields } \mathbf{v} \text{ s.t. } E_s(\mathbf{v}) = 0. \quad (5.4)$$

This condition is necessary for the well-posedness of problem (5.2)-(5.3). In the literature a tangential vector field  $\mathbf{v}$  defined on a surface and satisfying  $E_s(\mathbf{v}) = 0$  is known as a *Killing vector field* [39]. For a smooth two-dimensional Riemannian manifold, Killing vector fields form a Lie algebra of dimension at most 3. The subspace of all the Killing vector fields on  $\Gamma$  plays an important role in the analysis of problem (5.2)-(5.3).

For the weak formulation of problem (5.2)-(5.3), we use the spaces  $\mathbf{H}_t^1(\Gamma)$  and  $L_0^2(\Gamma) := \{p \in L^2(\Gamma) \mid \int_\Gamma p \, ds = 0\}$ . We also define the space of Killing vector fields

$$E := \{\mathbf{u} \in \mathbf{H}_t^1(\Gamma) \mid E_s(\mathbf{u}) = 0\}. \quad (5.5)$$

Note that  $E$  is a closed subspace of  $\mathbf{H}_t^1(\Gamma)$  and  $\dim(E) \leq 3$ .

Consider the bilinear forms (with  $A : B = \operatorname{tr}(AB^T)$  for  $A, B \in \mathbb{R}^{3 \times 3}$ )

$$a(\mathbf{u}, \mathbf{v}) := \int_\Gamma E_s(\mathbf{u}) : E_s(\mathbf{v}) \, ds, \quad \mathbf{u}, \mathbf{v} \in \mathbf{H}_t^1(\Gamma), \quad (5.6)$$

$$b(\mathbf{v}, p) := - \int_\Gamma p \operatorname{div}_\Gamma \mathbf{v} \, ds, \quad \mathbf{v} \in \mathbf{H}_t^1(\Gamma), \quad p \in L^2(\Gamma). \quad (5.7)$$

The weak (variational) formulation of the surface Stokes problem (5.2)-(5.3) reads: Determine  $(\mathbf{u}, p) \in \mathbf{H}_t^1(\Gamma)/E \times L_0^2(\Gamma)$  such that

$$\begin{aligned} a(\mathbf{u}, \mathbf{v}) + b(\mathbf{v}, p) &= (\mathbf{f}, \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathbf{H}_t^1(\Gamma)/E, \\ b(\mathbf{u}, q) &= 0 \quad \text{for all } q \in L^2(\Gamma). \end{aligned} \quad (5.8)$$

The following surface Korn inequality and inf-sup property were derived in [22].

LEMMA 5.1. *Assume  $\Gamma$  is  $C^2$  smooth and compact. There exist  $c_K > 0$  and  $c_0 > 0$  such that*

$$\|E_s(\mathbf{v})\|_{L^2(\Gamma)} \geq c_K \|\mathbf{v}\|_1 \quad \text{for all } \mathbf{v} \in \mathbf{H}_t^1(\Gamma)/E, \quad (5.9)$$

and

$$\sup_{\mathbf{v} \in \mathbf{H}_t^1(\Gamma)/E} \frac{b(\mathbf{v}, p)}{\|\mathbf{v}\|_1} \geq c_0 \|p\|_{L^2(\Gamma)} \quad \text{for all } p \in L_0^2(\Gamma). \quad (5.10)$$

Both bilinear forms  $a(\cdot, \cdot)$  and  $b(\cdot, \cdot)$  are also continuous. Therefore *problem (5.8) is well-posed*, and its unique solution is further denoted by  $\{\mathbf{u}^*, p^*\}$ .

We now introduce a stream function formulation. For this we need the following key assumption.

ASSUMPTION 5.1. *In the remainder we assume that  $\Gamma$  is simply connected.*

LEMMA 5.2. *The following relation holds for all  $\phi, \psi \in H^2(\Gamma)$ :*

$$\begin{aligned} a(\mathbf{curl}_\Gamma \phi, \mathbf{curl}_\Gamma \psi) &= \int_\Gamma E_s(\mathbf{curl}_\Gamma \phi) : E_s(\mathbf{curl}_\Gamma \psi) ds \\ &= \int_\Gamma \frac{1}{2} \Delta_\Gamma \phi \Delta_\Gamma \psi - K \nabla_\Gamma \phi \cdot \nabla_\Gamma \psi ds =: \tilde{a}(\phi, \psi). \end{aligned} \quad (5.11)$$

*Proof.* Since smooth functions are dense in  $H^2(\Gamma)$  it suffices to prove the relation for smooth functions  $\phi, \psi$ . Using partial integration and the identities in (2.5), (2.16), (2.12) we obtain

$$\begin{aligned} &\int_\Gamma E_s(\mathbf{curl}_\Gamma \phi) : E_s(\mathbf{curl}_\Gamma \psi) ds \\ &= \int_\Gamma \text{tr}(E_s(\mathbf{curl}_\Gamma \phi)(\nabla_\Gamma \mathbf{curl}_\Gamma \psi)) ds = - \int_\Gamma \mathbf{P} \text{div}_\Gamma(E_s(\mathbf{curl}_\Gamma \phi)) \cdot \mathbf{curl}_\Gamma \psi ds \\ &= -\frac{1}{2} \int_\Gamma [\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma \phi)) + 2K \mathbf{curl}_\Gamma \phi] \cdot \mathbf{curl}_\Gamma \psi ds \\ &= \frac{1}{2} \int_\Gamma (\mathbf{curl}_\Gamma \mathbf{curl}_\Gamma \phi)(\mathbf{curl}_\Gamma \mathbf{curl}_\Gamma \psi) - 2K \mathbf{curl}_\Gamma \phi \cdot \mathbf{curl}_\Gamma \psi ds \\ &= \int_\Gamma \frac{1}{2} \Delta_\Gamma \phi \Delta_\Gamma \psi - K \nabla_\Gamma \phi \cdot \nabla_\Gamma \psi ds, \end{aligned}$$

which proves the desired result.  $\square$

We introduce some further notation for stream function spaces:

$$\begin{aligned} H_*^2(\Gamma) &:= H^2(\Gamma) \cap H_*^1(\Gamma), \quad \tilde{E} := \{ \psi \in H_*^2(\Gamma) \mid \tilde{a}(\psi, \psi) = 0 \} \\ \mathbf{H}_{t,\text{div}}^1 &:= \{ \mathbf{u} \in \mathbf{H}_t^1(\Gamma) \mid \text{div}_\Gamma \mathbf{u} = 0 \}. \end{aligned}$$

LEMMA 5.3. *The following holds:*

$$\mathbf{curl}_\Gamma : H_*^2(\Gamma) \rightarrow \mathbf{H}_{t,\text{div}}^1 \quad \text{is an homeomorphism,} \quad (5.12)$$

$$\mathbf{curl}_\Gamma : \tilde{E} \rightarrow E \quad \text{is an homeomorphism.} \quad (5.13)$$

*Proof.* Take  $\psi \in H_*^2(\Gamma)$ . From  $\mathbf{curl}_\Gamma \psi = 0$  it follows that  $\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma \psi) = \Delta_\Gamma \psi = 0$  on  $\Gamma$ . Hence,  $\psi$  is a constant function on  $\Gamma$ . Using  $\int_\Gamma \psi \, ds = 0$  it follows that  $\psi$  equals the zero function. Thus  $\mathbf{curl}_\Gamma$  is injective on  $H_*^2(\Gamma)$ , hence also on  $\tilde{E} \subset H_*^2(\Gamma)$ . Take  $\mathbf{u} \in \mathbf{H}_{t,\text{div}}^1$ . From the Helmholtz decomposition it follows that there exist (unique)  $\psi, \phi \in H_*^1(\Gamma)$  such that  $\mathbf{u} = \nabla_\Gamma \psi + \mathbf{curl}_\Gamma \phi$ . From  $\text{div}_\Gamma \mathbf{u} = 0$  it follows that  $\psi = 0$ . Hence,  $\mathbf{u} = \mathbf{curl}_\Gamma \phi = \mathbf{n} \times \nabla_\Gamma \phi$ , which implies  $\mathbf{n} \times \mathbf{u} = -\nabla_\Gamma \phi$ . From  $\mathbf{u} \in \mathbf{H}_t^1(\Gamma)$  it follows that  $\phi \in H^2(\Gamma)$ . Hence we have surjectivity and  $\mathbf{curl}_\Gamma : H_*^2(\Gamma) \rightarrow \mathbf{H}_{t,\text{div}}^1$  is an isomorphism. From  $\|\mathbf{curl}_\Gamma \phi\|_1 \leq c\|\phi\|_{H^2(\Gamma)}$  it follows that this isomorphism is bounded and using the open mapping theorem it follows that the mapping is an homeomorphism. Using  $a(\mathbf{curl}_\Gamma \phi, \mathbf{curl}_\Gamma \phi) = \tilde{a}(\phi, \phi)$  one easily checks that  $\mathbf{curl}_\Gamma(\tilde{E}) = E$ .  $\square$

The unique solution  $\mathbf{u}^*$  of the weak formulation (5.8) is also the unique solution of the following problem: determine  $\mathbf{u} \in \mathbf{H}_{t,\text{div}}^1/E$  such that

$$a(\mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathbf{H}_{t,\text{div}}^1/E. \quad (5.14)$$

THEOREM 5.4. *Let  $\mathbf{u}^* \in \mathbf{H}_{t,\text{div}}^1/E$  be the unique solution of (5.8) (or (5.14)) and  $\phi^* \in H_*^1(\Gamma)$  the unique stream function such that  $\mathbf{u}^* = \mathbf{curl}_\Gamma \phi^*$ . This  $\phi^*$  is the unique solution of the following problem: determine  $\phi \in H_*^2(\Gamma)/\tilde{E}$  such that*

$$\tilde{a}(\phi, \psi) = (\mathbf{f}, \mathbf{curl}_\Gamma \psi)_{L^2(\Gamma)} \quad \text{for all } \psi \in H_*^2(\Gamma)/\tilde{E}. \quad (5.15)$$

Furthermore, if  $\Gamma$  is  $C^3$ , the estimate

$$\|\phi^*\|_{H^3(\Gamma)} \leq c\|\mathbf{f}\|_{L^2(\Gamma)} \quad (5.16)$$

holds, with a constant  $c$  independent of  $\mathbf{f} \in \mathbf{L}_t^2(\Gamma)$ .

*Proof.* The mapping  $\mathbf{curl}_\Gamma : H_*^2(\Gamma)/\tilde{E} \rightarrow \mathbf{H}_{t,\text{div}}^1/E$  is an isomorphism. This implies

$$a(\mathbf{u}^*, \mathbf{v}) = (\mathbf{f}, \mathbf{v})_{L^2(\Gamma)} \quad \text{for all } \mathbf{v} \in \mathbf{H}_{t,\text{div}}^1/E$$

iff

$$a(\mathbf{curl}_\Gamma \phi^*, \mathbf{curl}_\Gamma \psi) = (\mathbf{f}, \mathbf{curl}_\Gamma \psi)_{L^2(\Gamma)} \quad \text{for all } \psi \in H_*^2(\Gamma)/\tilde{E}$$

iff

$$\tilde{a}(\phi, \psi) = (\mathbf{f}, \mathbf{curl}_\Gamma \psi)_{L^2(\Gamma)} \quad \text{for all } \psi \in H_*^2(\Gamma)/\tilde{E}.$$

Due to  $\mathbf{n} \times \mathbf{u}^* = -\nabla_\Gamma \phi^*$  and the  $H^2$ -regularity of (5.14) we have

$$\|\phi^*\|_{H^3(\Gamma)} \leq c\|\nabla_\Gamma \phi^*\|_{H^2(\Gamma)} = c\|\mathbf{n} \times \mathbf{u}^*\|_{H^2(\Gamma)} \leq c\|\mathbf{f}\|_{L^2(\Gamma)},$$

with a constant  $c$  independent of  $\mathbf{f}$ .  $\square$

For the discretization of the problem in stream function formulation it is convenient

to reformulate the fourth order problem (5.15) as a coupled system of two second order problems. This reformulation is given in the following lemma.

LEMMA 5.5. *Consider the following problem: determine  $\phi \in H_*^1(\Gamma)/\tilde{E}$ ,  $\xi \in H^1(\Gamma)$  such that*

$$\int_{\Gamma} \frac{1}{2} \nabla_{\Gamma} \xi \cdot \nabla_{\Gamma} \psi + K \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \psi ds = -(\mathbf{f}, \mathbf{curl}_{\Gamma} \psi)_{L^2(\Gamma)} \quad \forall \psi \in H_*^1(\Gamma)/\tilde{E} \quad (5.17)$$

$$\int_{\Gamma} \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \eta + \xi \eta ds = 0 \quad \forall \eta \in H^1(\Gamma). \quad (5.18)$$

We assume that  $\Gamma$  is  $C^3$ . This problem has a unique solution given by  $\hat{\phi} = \phi^*$ ,  $\hat{\xi} = \Delta_{\Gamma} \phi^*$ , with  $\phi^*$  the unique solution of (5.15).

*Proof.* Let  $\phi^*$  the unique solution of (5.15) and define  $\hat{\phi} := \phi^*$ ,  $\hat{\xi} := \Delta_{\Gamma} \phi^*$ . Note that due to (5.16) we have  $\hat{\xi} \in H^1(\Gamma)$ . From  $\hat{\xi} = \Delta_{\Gamma} \phi^* = \Delta_{\Gamma} \hat{\phi}$  it follows that the pair  $(\hat{\phi}, \hat{\xi})$  satisfies (5.18). From  $\tilde{a}(\phi^*, \psi) = (\mathbf{f}, \mathbf{curl}_{\Gamma} \psi)_{L^2(\Gamma)}$  for all  $\psi \in H_*^2(\Gamma)/\tilde{E}$ , partial integration and a density argument it follows that the pair  $(\hat{\phi}, \hat{\xi})$  also satisfies (5.17). We now prove uniqueness. Let  $(\hat{\phi}_1, \hat{\xi}_1)$ ,  $(\hat{\phi}_2, \hat{\xi}_2)$  be two solution pairs and define  $e_{\phi} := \hat{\phi}_1 - \hat{\phi}_2 \in H_*^1(\Gamma)/\tilde{E}$ ,  $e_{\xi} := \hat{\xi}_1 - \hat{\xi}_2 \in H^1(\Gamma)$ . From (5.18) and  $H^2$ -regularity of the Laplace-Beltrami equation we get  $\Delta_{\Gamma} e_{\phi} = e_{\xi}$  and  $e_{\phi} \in H_*^2(\Gamma)$ . From (5.17) we obtain

$$\int_{\Gamma} \frac{1}{2} \nabla_{\Gamma} e_{\xi} \cdot \nabla_{\Gamma} \psi + K \nabla_{\Gamma} e_{\phi} \cdot \nabla_{\Gamma} \psi ds = 0 \quad \forall \psi \in H_*^1(\Gamma)/\tilde{E},$$

and thus

$$\int_{\Gamma} -\frac{1}{2} \Delta_{\Gamma} e_{\phi} \Delta_{\Gamma} \psi + K \nabla_{\Gamma} e_{\phi} \cdot \nabla_{\Gamma} \psi ds = 0 \quad \forall \psi \in H_*^2(\Gamma)/\tilde{E}.$$

Taking  $\psi = e_{\phi}$  this implies  $\tilde{a}(e_{\phi}, e_{\phi}) = 0$ . From the definition of the kernel space  $\tilde{E}$  it follows that  $e_{\phi} = 0$  must hold. Hence, also  $e_{\xi} = 0$ .  $\square$

REMARK 5.1. From the definition of the kernel space  $E$  and the compatibility assumption (5.4) it follows that the test space  $\mathbf{H}_{t,\text{div}}^1/E$  in (5.14) can be replaced by the larger space  $\mathbf{H}_{t,\text{div}}^1$ . Using this one may check that the test space  $H_*^2(\Gamma)/\tilde{E}$  in (5.15) can be replaced by the larger space  $H_*^2(\Gamma)$  and that the test space  $H_*^1(\Gamma)/\tilde{E}$  in (5.17) can be replaced by the larger space  $H_*^1(\Gamma)$  and even by the space  $H^1(\Gamma)$ . These larger test spaces are more convenient for a finite element discretization.

REMARK 5.2. In view of the finite element discretization introduced in section 6 we derive another characterization of the kernel  $\tilde{E} = \{ \phi \in H_*^2(\Gamma) \mid \tilde{a}(\phi, \phi) = 0 \}$ , which allows a more feasible representation of the trial space  $H_*^1(\Gamma)/\tilde{E}$  used in Lemma 5.5. Let  $P_*$  denote the orthogonal projection on  $L^2$ , i.e.,  $P_* \phi = \phi - \frac{1}{|\Gamma|} \int_{\Gamma} \phi ds$ . We then have  $H_*^1(\Gamma) = P_*(H^1(\Gamma))$  and, using  $\tilde{a}(\phi, \phi) = \tilde{a}(P_* \phi, P_* \phi)$ , we get  $\tilde{E} = P_*(\hat{E})$  with  $\hat{E} := \{ \phi \in H^2(\Gamma) \mid \tilde{a}(\phi, \phi) = 0 \}$ . Note that  $1 \in \hat{E}$ . Let  $P_{\hat{E}}$  be the  $L^2$ -projection on  $\hat{E}$ . Consider a ( $L^2$ -orthogonal) direct sum  $H^1(\Gamma) = \hat{E} \oplus (I - P_{\hat{E}})H^1(\Gamma)$ . We then have

$$H_*^1(\Gamma)/\tilde{E} \simeq P_*(I - P_{\hat{E}})H^1(\Gamma) = (I - P_{\hat{E}})H^1(\Gamma), \quad (5.19)$$

where in the last equality we used  $1 \in \hat{E}$ . Using the relation (5.11) we obtain  $\hat{E} = \{ \phi \in H^2(\Gamma) \mid \tilde{a}(\phi, \psi) = 0 \text{ for all } \psi \in H^2(\Gamma) \}$ . Using similar arguments as in the

proof of Lemma 5.5 one can then show that  $\phi \in \hat{E}$  iff there exists  $\xi \in H^1(\Gamma)$  such that the pair  $(\phi, \xi) \in H^1(\Gamma)^2$  is a solution of:

$$\begin{aligned} \int_{\Gamma} \frac{1}{2} \nabla_{\Gamma} \xi \cdot \nabla_{\Gamma} \psi + K \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \psi \, ds &= 0 \quad \forall \psi \in H^1(\Gamma) \\ \int_{\Gamma} \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \eta + \xi \eta \, ds &= 0 \quad \forall \eta \in H^1(\Gamma). \end{aligned} \quad (5.20)$$

Based on the two remarks above we propose the following reformulation of the coupled problem described in Lemma 5.5.

1. Let  $\hat{E}$  be the finite dimensional space spanned by the  $\phi$  component of the solutions of the coupled homogeneous problem (5.20).
2. Solve the coupled problem: Determine  $\phi, \xi \in H^1(\Gamma)$  such that

$$\int_{\Gamma} \frac{1}{2} \nabla_{\Gamma} \xi \cdot \nabla_{\Gamma} \psi + K \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \psi \, ds = -(\mathbf{f}, \mathbf{curl}_{\Gamma} \psi)_{L^2(\Gamma)} \quad \forall \psi \in H^1(\Gamma) \quad (5.21)$$

$$\int_{\Gamma} \nabla_{\Gamma} \phi \cdot \nabla_{\Gamma} \eta + \xi \eta \, ds = 0 \quad \forall \eta \in H^1(\Gamma). \quad (5.22)$$

A solution is denoted by  $\tilde{\phi}, \tilde{\xi}$ .

3. The unique solution  $\phi^*$  as in Lemma 5.5 is given by  $\phi^* = (I - P_{\hat{E}}) \tilde{\phi}$ . This is the solution for the quotient space  $(I - P_{\hat{E}}) H^1(\Gamma)$ , cf. (5.19).

REMARK 5.3. From the discussion above we see that, if the space of Killing fields as dimension  $> 0$ , this causes some technical difficulties. This is not due to the use of the stream function formulation of the Stokes problem. Very similar difficulties arise if the Stokes problem in the  $(\mathbf{u}, p)$  variables is considered. In a time-dependent (Navier-)Stokes problem these difficulties vanish. In one time step of an implicit time discretization one has to solve a generalized stationary Stokes problem with an additional zero order term. The spatial operator (in the space of divergence free velocities) then is of the form  $-\mathbf{P} \operatorname{div}_{\Gamma}(E_s(\cdot)) + cI$  with a strictly positive constant  $c$  (inverse proportional to the time step). This operator has a zero kernel.

REMARK 5.4. For the (very) special case of a constant curvature  $K$  (i.e., a sphere), the coupled system (5.21)-(5.22), can be decoupled by eliminating  $\phi$  from (5.21) using (5.22) (and similarly for (5.20)).

**6. Finite element discretization and numerical experiment.** For the discretization of the stream function formulation we apply a Galerkin finite element method to the three-step variational formulation described above. In this paper we only present one particular Galerkin approach and show results of a numerical experiment with this finite element method. We neither present an error analysis of the finite element method nor a comparison with other methods. A detailed study of different finite element discretizations, including error analysis and an accurate method for reconstruction of  $\mathbf{u} = \mathbf{curl}_{\Gamma} \phi$  from the finite element approximation of the stream function  $\phi$ , will be treated in a forthcoming paper.

One good option for the discretization of the scalar surface PDEs (5.21)-(5.22) is the SFEM developed by Dziuk and Elliott, cf. e.g. [12, 13]. This method is used

for the discretization of a stream function formulation in [29]. We use another approach, namely the trace finite element approach (TraceFEM) [31]. We use the latter method because of the availability of software in our group that provides an easy implementation of a TraceFE discretization of (5.21)-(5.22). We briefly describe the method.

Let  $\Omega \subset \mathbb{R}^3$  be a fixed polygonal domain that strictly contains  $\Gamma$ . We consider a family of shape regular tetrahedral triangulations  $\{\mathcal{T}_h\}_{h>0}$  of  $\Omega$ . The surface  $\Gamma$  is approximated by a piecewise planar approximation as follows. We assume that  $\Gamma$  is the zero level of a level set function  $\phi$  (not necessarily a signed distance function). Let  $I_h$  be the piecewise linear nodal interpolation operator on  $\mathcal{T}_h$ . We define  $\Gamma_h := \{x \in \Omega \mid (I_h\phi)(x) = 0\}$ . The subset of tetrahedra that have a nonzero intersection with  $\Gamma_h$  is collected in the set denoted by  $\mathcal{T}_h^\Gamma$ . On  $\mathcal{T}_h^\Gamma$  we use a standard finite element space of continuous functions that are piecewise linear. This so-called *outer finite element space* is denoted by  $V_h$ . The nodal basis functions in  $V_h$  are denoted by  $\{\phi_i^h\}_{1 \leq i \leq m}$ . The finite element isomorphism that maps coefficients to functions is denoted by  $J_h : \mathbb{R}^m \rightarrow V_h$ ,  $J_h \mathbf{x} = \sum_{i=1}^m x_i \phi_i^h$ . The *trace finite element space* is obtained by simply taking traces of functions in  $V_h$ , i.e.,  $V_h^\Gamma := \{(\phi_h)|_{\Gamma_h} \mid \phi_h \in V_h\} \subset H^1(\Gamma_h)$ .

The discretization of (5.21)-(5.22) is as follows: determine  $\phi_h, \xi_h \in V_h^\Gamma$  such that

$$\int_{\Gamma_h} \frac{1}{2} \nabla_{\Gamma_h} \xi_h \cdot \nabla_{\Gamma_h} \psi_h + K_h \nabla_{\Gamma_h} \phi_h \cdot \nabla_{\Gamma_h} \psi_h ds = -(\mathbf{f}^e, \mathbf{curl}_{\Gamma_h} \psi_h)_{L^2(\Gamma_h)} \quad \forall \psi_h \in V_h^\Gamma \quad (6.1)$$

$$\int_{\Gamma_h} \nabla_{\Gamma_h} \phi_h \cdot \nabla_{\Gamma_h} \eta_h + \xi_h \eta_h ds = 0 \quad \forall \eta_h \in V_h^\Gamma. \quad (6.2)$$

Here  $\mathbf{f}^e$  denotes an extension of  $\mathbf{f}$  and  $K_h$  an approximation of the Gauss curvature  $K$ .

We introduce mass and stiffness matrices for the matrix-vector representation of the discrete problem. Define, for  $1 \leq i, j, \leq m$ :

$$M_{ij} = \int_{\Gamma_h} \phi_i^h \phi_j^h ds, \quad A_{ij} = \int_{\Gamma_h} \nabla_{\Gamma_h} \phi_i^h \cdot \nabla_{\Gamma_h} \phi_j^h ds, \quad A_{ij}^K = \int_{\Gamma_h} K_h \nabla_{\Gamma_h} \phi_i^h \cdot \nabla_{\Gamma_h} \phi_j^h ds.$$

The matrix-vector problem corresponding to (6.1)-(6.2) is of the form

$$\mathcal{A} \mathbf{y} = \mathbf{c}, \quad \text{with } \mathcal{A} = \begin{pmatrix} 2\mathbf{A}^K & \mathbf{A} \\ \mathbf{A} & \mathbf{M} \end{pmatrix}. \quad (6.3)$$

Note that  $\mathbf{M}$ ,  $\mathbf{A}$  are symmetric positive semidefinite and  $\mathbf{A}^K$ ,  $\mathcal{A}$  are in general only symmetric. The matrix  $\mathcal{A}$  is (close to) singular due to the fact that the constant function and the kernel space  $\hat{E}$  are not factored out. For the TraceFEM there is a further issue related to poor conditioning of  $\mathcal{A}$  resulting from the fact that the traces of the outer nodal basis functions in general do not form a (well-conditioned) basis of the trace finite element space. This difficulty can be solved by using an appropriate stabilization, e.g. [18]. Here we want to keep the method as simple as possible and therefore do not consider any stabilization.

The discretization of the coupled system of second order problems described in Lemma 5.5 is based on the three-step procedure given above:

**1.** For computing an *approximation*  $\hat{E}_h$  of the space  $\hat{E}$  we proceed as follows. We determine the (at most) 5 eigenvalues of  $\mathcal{A}$  with smallest absolute value and determine (heuristically) how many of these are “close to zero”, in the sense that we expect

these eigenvalues to converge to zero if  $h \downarrow 0$ . Let this number be  $p$ ,  $1 \leq p \leq 4$ , and  $\mathbf{v}^{(j)} \in \mathbb{R}^{2m}$ ,  $1 \leq j \leq p$ , the corresponding (orthogonal) eigenvectors. We restrict to the first  $m$  entries in these vectors (corresponding to the first block row in (6.3)), and the resulting vectors are denoted by  $\mathbf{w}^{(j)} \in \mathbb{R}^m$ ,  $1 \leq j \leq p$ . The corresponding finite element functions  $\phi_h^{(j)} := J_h \mathbf{w}^{(j)}$  span the space  $\hat{E}_h$ . We determine an  $L^2$ -orthogonal basis of  $\hat{E}_h$ .

**2.** We determine a solution  $\mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2)$ ,  $\mathbf{y}_1, \mathbf{y}_2 \in \mathbb{R}^m$  of the (singular but consistent) linear system (6.3).

**3.** Using the orthogonal basis in  $\hat{E}_h$ , we determine  $\phi_h = (I - P_{\hat{E}_h})(J_h \mathbf{y}_1)$ , which is the finite element approximation of the solution  $\phi^*$ .

### Numerical experiment.

We consider an ellipsoid  $\Gamma \subset \Omega := [-2, 2]^3$  given by

$$\Gamma := \{\mathbf{x} \in \mathbb{R}^3 : x_1^2 + x_2^2 + \left(\frac{x_3}{1.5}\right)^2 = 1\}$$

Using MAPLE the Gauss curvature of  $\Gamma$  can be determined:

$$K = 5.0625 \cdot \frac{2.25x_1^2 + 2.25x_2^2 + x_3^2}{(5.0625x_1^2 + 5.0625x_2^2 + x_3^2)^2}.$$

We choose a smooth function

$$\phi_{sol}(x_1, x_2, x_3) := x_2^2 + \sin(x_1 x_3) + x_1 x_2 x_3.$$

This function has a nonzero intersection with the kernel space  $\hat{E}$ , i.e.,  $\phi_{sol} \neq (I - P_{\hat{E}})\phi_{sol}$ . Using MAPLE we determine the corresponding scalar function  $-\frac{1}{2}\Delta_\Gamma^2 \phi^{sol} - \text{div}_\Gamma(K \nabla_\Gamma \phi^{sol})$  which is used as right hand side function  $\text{curl}_\Gamma \mathbf{f}$  in (5.21), (6.1).

For the discretization we use a tetrahedral triangulation of  $\Omega$  constructed by starting from a uniform subdivision of  $\Omega$  into 8 tetrahedra and then applying uniform refinement. The mesh size on refinement level  $\ell$  is denoted by  $h_\ell$ . The surface approximation  $\Gamma_h$  and the trace finite element space are constructed as explained above. We follow the procedure with steps 1-3 outlined above. The 5 smallest eigenvalues of the matrix  $\mathcal{A}$  are given in Table 6.1.

	$\lambda_1$	$\lambda_2$	$\lambda_3$	$\lambda_4$	$\lambda_5$
$\ell = 1$	$-1.9 \cdot 10^{-16}$	$2.9 \cdot 10^{-3}$	$-1.2 \cdot 10^{-2}$	$-1.4 \cdot 10^{-2}$	$-4.2 \cdot 10^{-2}$
$\ell = 2$	$-9.9 \cdot 10^{-18}$	$3.7 \cdot 10^{-4}$	$-2.4 \cdot 10^{-3}$	$-2.5 \cdot 10^{-3}$	$-3.5 \cdot 10^{-3}$
$\ell = 3$	$-5.1 \cdot 10^{-18}$	$2.7 \cdot 10^{-5}$	$-3.5 \cdot 10^{-4}$	$-3.5 \cdot 10^{-4}$	$-3.5 \cdot 10^{-4}$
$k = 4$	$-1.7 \cdot 10^{-17}$	$1.7 \cdot 10^{-6}$	$-1.2 \cdot 10^{-4}$	$-1.2 \cdot 10^{-4}$	$-1.2 \cdot 10^{-4}$
$\ell = 5$	$-3.9 \cdot 10^{-18}$	$1.1 \cdot 10^{-7}$	$-1.6 \cdot 10^{-5}$	$-1.6 \cdot 10^{-5}$	$-1.6 \cdot 10^{-5}$

Table 6.1: Smallest eigenvalues of the matrix  $\mathcal{A}$ .

The eigenvalue  $\lambda_1$  is zero within machine accuracy (corresponds to the constant function). We observe a large gap between  $\lambda_2$  and the eigenvalues  $\lambda_i$ ,  $i \geq 3$ . We expect that this eigenvalue approximates a zero eigenvalue of the continuous problem, and based on this we take  $p = 2$  and  $\hat{E}_h$  the two-dimensional space as explained in step 2 above. The kernel function  $\phi_h^{(2)}$  corresponding to  $\lambda_2$  is illustrated in Figure 6.1. We note that also the eigenvalues  $\lambda_i$ ,  $i = 3, 4, 5$ , are quite small (for increasing refinement

level). This is due to the fact that in the trace finite element method we did not use any stabilization, which leads to very poor conditioning of the stiffness matrix. We solve the linear system (6.3) using a preconditioned MINRES method (with only diagonal preconditioning). The resulting solution is then projected, as explained in step 3 above, to eliminate the kernel components, resulting in the finite element approximation  $\phi_h$ , shown in Figure 6.1.




Fig. 6.1: Color graphs of kernel function  $\phi_h^{(2)}$  (left) and discrete solution  $\phi_h$  (right).

In the solution  $\phi_{sol}$  we factor out the kernel (approximation), i.e. we determine  $\phi_h^* := (I - P_{\hat{E}_h})\phi_{sol}$ . The errors in the approximation  $\phi_h \approx \phi_h^*$  are shown in Table 6.2. We observe the optimal orders of convergence.

$\ell$	$\ \phi_h - \phi_h^*\ _{L^2(\Gamma_h)}$	$EOC$	$ \phi_h - \psi_h^* _{H^1(\Gamma_h)}$	$EOC$
1	$6.63 \cdot 10^{-1}$		$3.27 \cdot 10^0$	
2	$2.04 \cdot 10^{-1}$	1.70	$1.57 \cdot 10^0$	1.06
3	$5.81 \cdot 10^{-2}$	1.81	$7.62 \cdot 10^{-1}$	1.04
4	$1.50 \cdot 10^{-2}$	1.95	$3.80 \cdot 10^{-1}$	1.00
5	$3.67 \cdot 10^{-3}$	2.03	$1.90 \cdot 10^{-1}$	1.00

Table 6.2: Discretization errors

**7. Appendix.** In this section we derive the results (2.7), (2.15) and (2.13). The proofs are based on elementary tensor calculus. We use standard tensor notation and the Einstein summation convention (always over  $i = 1, 2, 3$ , for repeated indices  $i$ ). For a scalar function  $\phi$  we have, cf. (2.1):

$$(\nabla_{\Gamma}\phi)_i = P_{ik}\partial_k\phi.$$

(scalar entries of the matrix  $\mathbf{P}$  are denoted  $P_{ij}$ ). For the vector function  $\mathbf{u} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  we have, cf. (2.2), (2.3):

$$(\nabla_{\Gamma}\mathbf{u})_{ij} = P_{ik}\partial_l u_k P_{lj}, \quad \operatorname{div}_{\Gamma}\mathbf{u} = (\nabla_{\Gamma}\mathbf{u})_{ii} = P_{ik}\partial_k u_l P_{li} = P_{lk}\partial_k u_l,$$



and for the matrix divergence operator (2.3) we have the representation:

$$(\operatorname{div}_\Gamma A)_i = \operatorname{div}_\Gamma(e_i^T A) = P_{lk}\partial_k A_{il}. \quad (7.1)$$

For manipulations of vector products it is convenient to use the three-dimensional Levi-Civita symbol (also called permutation tensor):

$$\epsilon_{ijk} := \begin{cases} +1 & \text{if } (ijk) \text{ is an even permutation of } (123) \\ -1 & \text{if } (ijk) \text{ is an odd permutation of } (123) \\ 0 & \text{otherwise.} \end{cases}$$

This tensor is antisymmetric, e.g.,  $\epsilon_{ijk} = -\epsilon_{jik}$  for all  $i, j, k$ . For vectors  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$  we have  $(\mathbf{a} \times \mathbf{b})_k = \epsilon_{ijk} a_i b_j$ ,  $k = 1, 2, 3$ . We will also use the relation

$$\epsilon_{jki}\epsilon_{nml} = \delta_{jn}\delta_{km}\delta_{il} - \delta_{jn}\delta_{kl}\delta_{im} - \delta_{jm}\delta_{kn}\delta_{il} + \delta_{jm}\delta_{kl}\delta_{in} + \delta_{jl}\delta_{kn}\delta_{im} - \delta_{jl}\delta_{km}\delta_{in}, \quad (7.2)$$

with  $\delta_{ij}$  the Kronecker symbol, i.e.,  $\delta_{ii} = 1$ , and zero otherwise. We will often use the following relations, with the symmetric Weingarten mapping denoted by  $\mathbf{H} = \nabla \mathbf{n}$ , which satisfies  $\mathbf{P}\mathbf{H} = \mathbf{H} = \mathbf{H}\mathbf{P}$ ,  $\mathbf{H}\mathbf{n} = 0$ , and the notation  $\partial_k = \partial_{x_k}$  for the  $k$ -th partial derivative in  $\mathbb{R}^3$ :

$$P_{ik}n_k = 0, \quad H_{ik}n_k = 0, \quad \partial_k P_{ij} = -n_i H_{kj} - n_j H_{ki}. \quad (7.3)$$

We first derive the identity (2.7). Note that

$$(\nabla_\Gamma \times \mathbf{u}) \cdot \mathbf{n} = \epsilon_{ijl} P_{ik} \partial_k u_j n_l. \quad (7.4)$$

We also have

$$\begin{aligned} \operatorname{div}_\Gamma(\mathbf{u} \times \mathbf{u}) &= P_{ik} \partial_k (\mathbf{u} \times \mathbf{n})_i = P_{ik} \partial_k (\epsilon_{jli} u_j n_l) = \epsilon_{jli} P_{ik} \partial_k (u_j n_l) \\ &= \epsilon_{jli} P_{ik} \partial_k u_j n_l + \epsilon_{jli} P_{ik} u_j H_{kl}. \end{aligned} \quad (7.5)$$

For the last term we get  $\epsilon_{jli} P_{ik} u_j H_{kl} = \epsilon_{jli} u_j H_{il}$ . Using the antisymmetry property of the Levi-Civita symbol and the symmetry of  $\mathbf{H}$  we get  $\epsilon_{jli} H_{il} = -\epsilon_{jil} H_{li} = -\epsilon_{jli} H_{il}$ , hence,  $\epsilon_{jli} u_j H_{il} = 0$ , i.e., the last term in (7.5) vanishes. Using the permutation properties of the Levi-Civita symbol we get  $\epsilon_{jli} = \epsilon_{ijl}$  and using this in (7.5) and comparing with (7.4) yields the relation (2.7).

We derive the result (2.15). Using the representations and relations introduced above we get

$$\begin{aligned} \operatorname{div}_\Gamma(\mathbf{n} \times \nabla_\Gamma \phi) &= P_{lk} \partial_k (\mathbf{n} \times \nabla_\Gamma \phi)_l = P_{lk} \partial_k (\epsilon_{ijl} n_i P_{jr} \partial_r \phi) \\ &= \epsilon_{ijl} P_{lk} (H_{ki} P_{jr} \partial_r \phi + n_i \partial_k P_{jr} \partial_r \phi + n_i P_{jr} \partial_k \partial_r \phi) \\ &= \epsilon_{ijl} (H_{li} P_{jr} \partial_r \phi - n_i n_j H_{lr} \partial_r \phi - n_i n_r H_{lj} \partial_r \phi + n_i P_{lk} P_{jr} \partial_k \partial_r \phi). \end{aligned}$$

Using the antisymmetry property of the Levi-Civita symbol and the symmetry of  $\mathbf{H}$  we get  $\epsilon_{ijl} H_{li} = 0$ , hence  $\epsilon_{ijl} H_{li} P_{jr} \partial_r \phi = 0$ . The other three terms can be treated similarly, since  $n_i n_j$  is symmetric w.r.t.  $(ij)$ ,  $H_{lj}$  is symmetric w.r.t.  $(jl)$  and  $P_{lk} P_{jr} \partial_k \partial_r \phi$  is symmetric w.r.t.  $(jl)$ . From this it follows that  $\operatorname{div}_\Gamma(\mathbf{n} \times \nabla_\Gamma \phi) = 0$ .

The proof of (2.13) requires a more tedious derivation. From the definitions and representations given above we get, using  $\epsilon_{nml} H_{ln} = 0$  (due to antisymmetry),

$$\begin{aligned} \operatorname{curl}_\Gamma \mathbf{u} &= \operatorname{div}_\Gamma(\mathbf{u} \times \mathbf{n}) = -P_{lr} \partial_r (\mathbf{n} \times \mathbf{u})_l = -P_{lr} \partial_r (\epsilon_{nml} n_n u_m) \\ &= -\epsilon_{nml} P_{lr} (H_{rn} u_m + n_n \partial_r u_m) = -\epsilon_{nml} H_{ln} u_m - \epsilon_{nml} P_{lr} n_n \partial_r u_m \\ &= -\epsilon_{nml} P_{lr} n_n \partial_r u_m. \end{aligned}$$

Using this we obtain

$$\begin{aligned} (\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma \mathbf{u}))_i &= (\mathbf{n} \times \nabla_\Gamma(\mathbf{curl}_\Gamma \mathbf{u}))_i = \epsilon_{jki} n_j (\nabla_\Gamma \mathbf{curl}_\Gamma \mathbf{u})_k = \epsilon_{jki} n_j P_{ks} \partial_s (\mathbf{curl}_\Gamma \mathbf{u}) \\ &= -\epsilon_{jki} n_j P_{ks} \partial_s (\epsilon_{nml} P_{lr} n_n \partial_r u_m) = -\epsilon_{jki} \epsilon_{nml} n_j P_{ks} \partial_s (P_{lr} n_n \partial_r u_m) \end{aligned}$$

We now use the identity (7.2), which results in 6 nonzero terms, namely for  $(nml) \in \{(jki), (jik), (kji), (ijk), (kij), (ikj)\}$  with a corresponding sign as in (7.2). This yields

$$\begin{aligned} (\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma \mathbf{u}))_i &= -n_j P_{ks} \partial_s (P_{ir} n_j \partial_r u_k) + n_j P_{ks} \partial_s (P_{kr} n_j \partial_r u_i) \\ &\quad + n_j P_{ks} \partial_s (P_{ir} n_k \partial_r u_j) - n_j P_{ks} \partial_s (P_{ks} n_i \partial_r u_j) \\ &\quad - n_j P_{ks} \partial_s (P_{jr} n_k \partial_r u_i) + n_j P_{ks} \partial_s (P_{jr} n_i \partial_r u_k) \\ &=: (1) + (2) + (3) + (4) + (5) + (6). \end{aligned} \tag{7.6}$$

We now analyze these 6 terms. We start with the fifth one. Using  $\mathbf{Pn} = 0$  we get

$$(5) = -n_j P_{ks} \partial_s (P_{jr} n_k \partial_r u_i) = -n_j P_{ks} P_{jr} H_{sk} \partial_r u_i = 0. \tag{7.7}$$

For the third term we get

$$(3) = n_j P_{ks} \partial_s (P_{ir} n_k \partial_r u_j) = n_j P_{ks} P_{ir} H_{sk} \partial_r u_j = n_j H_{ss} P_{ir} \partial_r u_j. \tag{7.8}$$

We take the first and sixth term together:

$$\begin{aligned} (1) + (6) &= n_j P_{ks} \partial_s \left( (P_{jr} n_i - P_{ir} n_j) \partial_r u_k \right) \\ &= n_j P_{ks} \partial_s (P_{jr} n_i - P_{ir} n_j) \partial_r u_k - P_{ks} P_{ir} \partial_s \partial_r u_k. \end{aligned}$$

Now note (we use (7.3)):

$$\begin{aligned} n_j P_{ks} \partial_s (P_{jr} n_i - P_{ir} n_j) &= P_{ks} \partial_s \left( n_j (P_{jr} n_i - P_{ir} n_j) \right) - P_{ks} H_{sj} (P_{jr} n_i - P_{ir} n_j) \\ &= -P_{ks} \partial_s P_{ir} - P_{ks} H_{sr} n_i \\ &= P_{ks} (n_i H_{sr} + n_r H_{si}) - P_{ks} H_{sr} n_i = n_r H_{ki}. \end{aligned}$$

Hence,

$$(1) + (6) = n_r H_{ki} \partial_r u_k - P_{ks} P_{ir} \partial_s \partial_r u_k. \tag{7.9}$$

Finally we combine the second and fourth term. We use  $P_{ks} \partial_s P_{kr} = \partial_s P_{sr} = -n_r H_{ss}$ ,  $n_r n_j \partial_r u_j = 0$  (which follows from  $\partial_r (n_j u_j) = 0$  and  $n_r H_{rj} = 0$ ) and  $n_r \partial_r u_i = n_r \partial_r (P_{ki} u_k) = n_r P_{ki} \partial_r u_k$ , and then get:

$$\begin{aligned} (2) + (4) &= n_j P_{ks} \partial_s \left( P_{kr} (n_j \partial_r u_i - n_i \partial_r u_j) \right) \\ &= -n_j n_r H_{ss} (n_j \partial_r u_i - n_i \partial_r u_j) + n_j P_{sr} \partial_s (n_j \partial_r u_i - n_i \partial_r u_j) \\ &= -n_r H_{ss} \partial_r u_i + n_j P_{sr} \left( H_{sj} \partial_r u_i + n_j \partial_s \partial_r u_i - H_{si} \partial_r u_j - n_i \partial_s \partial_r u_j \right) \\ &= -n_r H_{ss} P_{ki} \partial_r u_k + P_{sr} \partial_s \partial_r u_i - n_j H_{ri} \partial_r u_j - n_i n_j P_{sr} \partial_s \partial_r u_j. \end{aligned}$$

Note that  $-n_i n_j P_{sr} \partial_s \partial_r u_j = P_{ij} P_{sr} \partial_s \partial_r u_j - P_{sr} \partial_s \partial_r u_i$ . Hence we get

$$(2) + (4) = -n_r H_{ss} P_{ki} \partial_r u_k - n_k H_{ri} \partial_r u_k + P_{ij} P_{sr} \partial_s \partial_r u_j. \tag{7.10}$$

Substitution of the results in (7.7)-(7.10) in (7.6) yields:

$$\begin{aligned}
(\mathbf{curl}_\Gamma(\mathbf{curl}_\Gamma \mathbf{u}))_i &= n_k H_{ss} P_{ir} \partial_r u_k + n_r H_{ki} \partial_r u_k - P_{ks} P_{ir} \partial_s \partial_r u_k \\
&\quad - n_r H_{ss} P_{ki} \partial_r u_k - n_k H_{ri} \partial_r u_k + P_{ik} P_{sr} \partial_s \partial_r u_k \\
&= [n_r H_{ik} - n_k H_{ir} + H_{ss} (n_k P_{ir} - n_r P_{ik})] \partial_r u_k \\
&\quad + (P_{ik} P_{sr} - P_{ir} P_{sk}) \partial_s \partial_r u_k.
\end{aligned} \tag{7.11}$$

We now consider the expression on the right hand side in (2.13). Note that

$$(\nabla_\Gamma \mathbf{u} - \nabla_\Gamma \mathbf{u}^T)_{nm} = P_{nk} P_{rm} \partial_r u_k - P_{mk} P_{rn} \partial_r u_k = (P_{mr} P_{kn} - P_{nr} P_{km}) \partial_r u_k$$

Using (7.1) we get  $(\mathbf{P} \operatorname{div}_\Gamma A)_i = P_{in} P_{ms} \partial_s A_{nm}$  and thus

$$\begin{aligned}
(\mathbf{P} \operatorname{div}_\Gamma (\nabla_\Gamma \mathbf{u} - \nabla_\Gamma \mathbf{u}^T))_i &= P_{in} P_{ms} \partial_s ((P_{mr} P_{kn} - P_{nr} P_{km}) \partial_r u_k) \\
&= P_{in} P_{ms} (P_{mr} P_{kn} - P_{nr} P_{km}) \partial_s \partial_r u_k + P_{in} P_{ms} \partial_s (P_{mr} P_{kn} - P_{nr} P_{km}) \partial_r u_k \\
&= (P_{ik} P_{sr} - P_{ir} P_{sk}) \partial_s \partial_r u_k + P_{in} P_{ms} \partial_s (P_{mr} P_{kn} - P_{nr} P_{km}) \partial_r u_k.
\end{aligned}$$

We also have

$$\begin{aligned}
&P_{in} P_{ms} \partial_s (P_{mr} P_{kn} - P_{nr} P_{km}) \\
&= P_{in} P_{ms} [(-n_m H_{sr} - n_r H_{sm}) P_{kn} + (-n_k H_{sn} - n_n H_{sk}) P_{mr} \\
&\quad + (n_n H_{sr} + n_r H_{sn}) P_{km} + (n_k H_{sm} + n_m H_{sk}) P_{nr}] \\
&= -n_r H_{ss} P_{ik} - n_k H_{ir} + n_r H_{ik} + n_k H_{ss} P_{ir}
\end{aligned}$$

Combination of the above two results yields

$$\begin{aligned}
(\mathbf{P} \operatorname{div}_\Gamma (\nabla_\Gamma \mathbf{u} - \nabla_\Gamma \mathbf{u}^T))_i &= (P_{ik} P_{sr} - P_{ir} P_{sk}) \partial_s \partial_r u_k \\
&\quad + [n_r H_{ik} - n_k H_{ir} + H_{ss} (n_k P_{ir} - n_r P_{ik})] \partial_r u_k
\end{aligned}$$

and comparing this with (7.11) completes the proof of (2.13).

**Acknowledgements** The fruitful discussions on the topic of this paper with Philip Brandner and Thomas Jankuhn are acknowledged.

#### REFERENCES

- [1] R. ABRAHAM, J. MARSDEN, AND T. RATIU, *Manifolds, Tensor Analysis, and Applications*, Springer, 1988.
- [2] M. ARNAUDON AND A. B. CRUZEIRO, *Lagrangian Navier–Stokes diffusions on manifolds: variational principle and stability*, Bulletin des Sciences Mathématiques, 136 (2012), pp. 857–881.
- [3] V. I. ARNOLD, *Mathematical Methods of Classical Mechanics*, vol. 60, Springer Science & Business Media, 2013.
- [4] M. ARROYO AND A. DESIMONE, *Relaxation dynamics of fluid membranes*, Physical Review E, 79 (2009), p. 031915.
- [5] J. W. BARRETT, H. GARCKE, AND R. NÜRNBERG, *A stable numerical method for the dynamics of fluidic membranes*, Numerische Mathematik, 134 (2016), pp. 783–822.
- [6] R. BOTT AND L. W. TU, *Differential Forms in Algebraic Topology*, vol. 82 of Graduate Texts in Mathematics, Springer, 1982.
- [7] H. BRENNER, *Interfacial Transport Processes and Rheology*, Elsevier, 2013.
- [8] A. BUFFA AND P. CHARLET JR., *On traces for functional spaces related to Maxwell’s equations Part II: Hodge decompositions on the boundary of Lipschitz polyhedra and applications*, Math. Methods Appl. Sci., 24 (2001), pp. 31–48.

- [9] M. CESSENAT, *Mathematical Methods in Electromagnetism*, World Scientific Publishing Co., 1996.
- [10] C. DIERECK AND F. CROWET, *Helmholtz decomposition on multiply connected domains*, Philips J. Res., 39 (1984), pp. 242–253.
- [11] M. DO CARMO, *Differential Geometry of Curves and Surfaces*, Prentice-Hall, 1976.
- [12] G. DZIUK AND C. ELLIOTT, *Finite elements on evolving surfaces*, IMA J. Numer. Anal., 27 (2007), pp. 262–292.
- [13] G. DZIUK AND C. M. ELLIOTT, *Finite element methods for surface PDEs*, Acta Numerica, 22 (2013), pp. 289–396.
- [14] D. G. EBIN AND J. MARSDEN, *Groups of diffeomorphisms and the motion of an incompressible fluid*, Annals of Mathematics, (1970), pp. 102–163.
- [15] P. FIRBY AND C. GARDINER, *Surface Topology*, Ellis Horwood Limited, 1982.
- [16] T.-P. FRIES, *Higher-order surface FEM for incompressible Navier-Stokes flows on manifolds*, arXiv:1712.02520, (2017).
- [17] V. GIRAULT AND P. A. RAVIART, *Finite Element Methods for Navier-Stokes Equations*, Springer, Berlin, 1986.
- [18] J. GRANDE AND A. REUSKEN, *A higher order finite element method for partial differential equations on surfaces*, SIAM Journal on Numerical Analysis, 54 (2016), pp. 388–414.
- [19] M. E. GURTIN AND A. I. MURDOCH, *A continuum theory of elastic material surfaces*, Archive for Rational Mechanics and Analysis, 57 (1975), pp. 291–323.
- [20] P. HANSBO AND M. LARSON, *Continuous/discontinuous finite element modelling of Kirchhoff plate structures in  $r^3$  using tangential differential calculus*, Computational Mechanics, 60 (2017), pp. 693–702.
- [21] P. HANSBO, M. G. LARSON, AND K. LARSSON, *Analysis of finite element methods for vector Laplacians on surfaces*, arXiv preprint arXiv:1610.06747, (2016).
- [22] T. JANKUHN, M. OLSHANSKII, AND A. REUSKEN, *Incompressible fluid problems on embedded surfaces: Modeling and variational formulations*, preprint 462, IGPM, RWTH Aachen University, arXiv:1702.02989, 2017.
- [23] H. Koba, C. LIU, AND Y. GIGA, *Energetic variational approaches for incompressible fluid systems on an evolving surface*, Quarterly of Applied Mathematics, (2016).
- [24] W. KÜHNEL, *Differential Geometry: Curves-Surfaces-Manifolds*, American Mathematical Society, 2015.
- [25] I. MADSEN AND J. TORNEHAVE, *From Calculus to Cohomology: de Rham Cohomology and Characteristic Classes*, Cambridge University Press, 1997.
- [26] M. MITREA AND M. TAYLOR, *Navier-Stokes equations on Lipschitz domains in Riemannian manifolds*, Mathematische Annalen, 321 (2001), pp. 955–987.
- [27] T.-H. MIURA, *On singular limit equations for incompressible fluids in moving thin domains*, arXiv:1703.09698, (2017).
- [28] C. B. MORREY, *Multiple Integrals in the Calculus of Variations*, no. 130 in Die Grundlagen der mathematischen Wissenschaften in Einzeldarstellungen, Springer, 1966.
- [29] I. NITSCHKE, A. VOIGT, AND J. WENSCH, *A finite element approach to incompressible two-phase flow on manifolds*, Journal of Fluid Mechanics, 708 (2012), pp. 418–438.
- [30] M. OLSHANSKII, A. QUAINI, A. REUSKEN, AND V. YUSHUTIN, *A finite element method for the surface Stokes problem*, preprint 475, arxiv:1801.06589, IGPM, RWTH Aachen University, 2017.
- [31] M. OLSHANSKII, A. REUSKEN, AND J. GRANDE, *A finite element method for elliptic equations on surfaces*, SIAM J. Numer. Anal., 47 (2009), pp. 3339–3358.
- [32] A. QUARTERONI AND A. VALLI, *Numerical Approximation of Partial Differential Equations*, Springer, Berlin, 1994.
- [33] M. RAHIMI, A. DESIMONE, AND M. ARROYO, *Curved fluid membranes behave laterally as effective viscoelastic media*, Soft Matter, 9 (2013), pp. 11033–11045.
- [34] P. RANGAMANI, A. AGRAWAL, K. K. MANDADAPU, G. OSTER, AND D. J. STEIGMANN, *Interaction between surface shape and intra-surface viscous flow on lipid membranes*, Biomechanics and modeling in mechanobiology, (2013), pp. 1–13.
- [35] A. REUSKEN AND Y. ZHANG, *Numerical simulation of incompressible two-phase flows with a Boussinesq-Scriven surface stress tensor*, Numerical Methods in Fluids, 73 (2013), pp. 1042–1058.
- [36] S. REUTHER AND A. VOIGT, *The interplay of curvature and vortices in flow on curved surfaces*, Multiscale Modeling & Simulation, 13 (2015), pp. 632–643.
- [37] ———, *Solving the incompressible surface Navier-Stokes equation by surface finite elements*, arXiv preprint arXiv:1709.02803, (2017).
- [38] G. S., T. JANKUHN, M. OLSHANSKII, AND A. REUSKEN, *A trace finite element method for vector-*

- Laplacians on surfaces*, preprint 469, IGPM, RWTH Aachen University, arXiv:1709.00479, 2017.
- [39] T. SAKAI, *Riemannian geometry*, vol. 149, American Mathematical Soc., 1996.
  - [40] L. SCRIVEN, *Dynamics of a fluid interface equation of motion for Newtonian surface fluids*, *Chemical Engineering Science*, 12 (1960), pp. 98–108.
  - [41] J. C. SLATTERY, L. SAGIS, AND E.-S. OH, *Interfacial transport phenomena*, Springer Science & Business Media, 2007.
  - [42] L. TARTAR, *Nonlinear differential equations using compactness methods*, Report 1584, Mathematics Research Center, Univ. of Wisconsin, Madison, 1975.
  - [43] M. E. TAYLOR, *Analysis on Morrey spaces and applications to Navier-Stokes and other evolution equations*, *Communications in Partial Differential Equations*, 17 (1992), pp. 1407–1456.
  - [44] R. TEMAM, *Infinite-dimensional dynamical systems in mechanics and physics*, Springer, New York, 1988.
  - [45] J. THORPE, *Elementary Topics in Differential Geometry*, Undergraduate Texts in Mathematics, Springer, 1997.
  - [46] J. WLOKA, *Partial Differential Equations*, Cambridge University Press, Cambridge, 1987.