

ANALYSIS OF A STOKES INTERFACE PROBLEM

MAXIM A. OLSHANSKII* AND ARNOLD REUSKEN †

Abstract. We consider a stationary Stokes problem with a piecewise constant viscosity coefficient. For the variational formulation of this problem we prove a well-posedness result in which the constants are uniform with respect to the jump in the viscosity coefficient. We apply a standard discretization with a pair of LBB stable finite element spaces. The main result of the paper is an infsup result for the discrete problem that is uniform with respect to the jump in the viscosity coefficient. From this we derive a robust estimate for the discretization error. We prove that the mass matrix with respect to some suitable scalar product yields a robust preconditioner for the Schur complement. Results of numerical experiments are presented that illustrate this robustness property.

AMS subject classifications. 65N15, 65N22, 65N30, 65F10

Key words. Stokes equations, interface problem, infsup condition, finite elements, preconditioning, Schur complement

1. Introduction. In this paper we treat the following Stokes problem on a bounded connected Lipschitz domain Ω in d -dimensional Euclidean space ($d = 2, 3$): Find a velocity \mathbf{u} and a pressure p such that

$$-\operatorname{div}(\nu(\mathbf{x})\nabla\mathbf{u}) + \nabla p = \mathbf{f} \quad \text{in } \Omega, \quad (1.1)$$

$$\operatorname{div} \mathbf{u} = 0 \quad \text{in } \Omega, \quad (1.2)$$

$$\mathbf{u} = 0 \quad \text{on } \partial\Omega, \quad (1.3)$$

with a *piecewise constant viscosity*:

$$\nu = \begin{cases} 1 & \text{in } \Omega_1 \\ \varepsilon > 0 & \text{in } \Omega_2. \end{cases} \quad (1.4)$$

The subdomains Ω_1, Ω_2 are assumed to be Lipschitz domains such that $\Omega_1 \cap \Omega_2 = \emptyset$ and $\overline{\Omega} = \overline{\Omega}_1 \cup \overline{\Omega}_2$. By Γ we denote the interface between the subdomains: $\Gamma = \partial\Omega_1 \cap \partial\Omega_2$.

An important motivation for considering this type of Stokes equations comes from two-phase incompressible flows. Often such problems are modeled by Navier-Stokes equations with discontinuous density and viscosity coefficients. The effect of interface tension can be taken into account by using a special localized force term at the interface. The latter approach is known as the continuum surface force (CSF) model, cf. [5]. A well-known technique for capturing the unknown interface is based on the level set method, cf. [26, 20, 14] and the references therein. If in such a setting one has highly viscous flows then the Stokes equations with discontinuous viscosity are a reasonable model problem. We note that from a modeling point of view it would be better to take the full velocity tensor $D(\mathbf{u}) := \frac{1}{2}(\nabla\mathbf{u} + (\nabla\mathbf{u})^T)$ instead of $\nabla\mathbf{u}$ in (1.1). To make the presentation more transparent we decided to present the analysis for the case with $\nabla\mathbf{u}$. The analysis can be extended to the case with the full velocity tensor. We comment on this in section 6.

* Department of Mechanics and Mathematics, Moscow State University, Moscow 119899, Russia; email: Maxim.Olshanskii@mtu-net.ru. This author was supported by the German Research Foundation through the guest program of SFB 540

† Institut für Geometrie und Praktische Mathematik, RWTH-Aachen, D-52056 Aachen, Germany; email: reusken@igpm.rwth-aachen.de

For pure diffusion problems (Poisson equation) with a discontinuous diffusion coefficient one can find analyses of discretization methods [1, 2, 6, 15, 22], error estimators [21, 4] and iterative solvers [8, 9, 23, 28] in the literature. For the Stokes problem with discontinuous viscosity, however, we did not find any theoretical analysis focusing on ν -dependence of results. This paper presents an analysis of a finite element method and of a solver for the discretized Stokes interface problem together with some results for the weak formulation of (1.1)-(1.3).

We introduce a variational formulation of the equations (1.1)-(1.3). We use the notation $\mathbf{V} := H_0^1(\Omega)^d$ for the velocity space. For the pressure space some factorization of $L^2(\Omega)$ is used. It appears that for this problem it is convenient to use:

$$M := \{ p \in L^2(\Omega) \mid \int_{\Omega} \nu^{-1} p(x) dx = 0 \} . \quad (1.5)$$

The variational problem reads as follows: given $\mathbf{f} \in \mathbf{V}'$ find $\{\mathbf{u}, p\} \in \mathbf{V} \times M$ such that

$$\begin{cases} (\nu \nabla \mathbf{u}, \nabla \mathbf{v}) - (\operatorname{div} \mathbf{v}, p) = (\mathbf{f}, \mathbf{v}) & \text{for } \mathbf{v} \in \mathbf{V}, \\ (\operatorname{div} \mathbf{u}, q) = 0 & \text{for } q \in M . \end{cases} \quad (1.6)$$

Here and in the remainder the L^2 scalar product and associated norm are denoted by (\cdot, \cdot) , $\|\cdot\|$, respectively. The bilinear form $(\nu \nabla \cdot, \nabla \cdot)$ defines a scalar product on \mathbf{V} . We use the norm induced by this scalar product:

$$\|\mathbf{u}\|_{\mathbf{V}} := (\nu \nabla \mathbf{u}, \nabla \mathbf{u})^{\frac{1}{2}} \quad \text{for } \mathbf{u} \in \mathbf{V} \quad (1.7)$$

On M , apart from the L^2 scalar product we will also use a weighted L^2 scalar product:

$$(p, q)_M := \int_{\Omega} \nu^{-1} p q dx = (\nu^{-1} p, q) \quad \text{for } p, q \in M , \quad (1.8)$$

and $\|p\|_M := (p, p)_M^{\frac{1}{2}}$. In the analysis we use the ν -dependent norm $(\|\cdot\|_{\mathbf{V}}^2 + \|\cdot\|_M^2)^{\frac{1}{2}}$ on the product space $\mathbf{V} \times M$. In section 2 we prove a continuity and an infsup result that are uniform with respect to the parameter ε . Using standard arguments this then yields uniform well-posedness of the continuous Stokes problem.

In section 3 we consider the discrete variational problem in a pair of finite element spaces $(M_h \subset M, \mathbf{V}_h \subset \mathbf{V})$ that are assumed to be LBB stable. As a main result of this paper we present a discrete infsup result that is uniform with respect to the parameters h (mesh size) and ε . This result is used to derive a (sharp) uniform bound for the discretization error.

In section 4 we prove that the mass-matrix with respect to the scalar product (1.8) in the pressure subspace M_h is spectrally equivalent to the Schur complement *uniform in the parameters h and ε* . In combination with known results on block-preconditioning and on multigrid this then implies optimality results for certain iterative methods. For the Uzawa method and a preconditioned MINRES method we present results of numerical experiments in section 5. In section 6 we comment on closely related Stokes interface problems. We consider the case with the full velocity tensor $D(\mathbf{u})$ instead of $\nabla \mathbf{u}$ in (1.1) and the problem where instead of M we use the more standard space $L_0^2(\Omega)$ for the pressure.

2. The continuous problem. In this section we analyze the variational problem (1.6). By definition we have ellipticity and continuity of the bilinear form $(\nu \nabla \cdot, \nabla \cdot)$

in the space $(\mathbf{V}, \|\cdot\|_{\mathbf{V}}) : (\nu \nabla \mathbf{u}, \nabla \mathbf{u}) = \|\mathbf{u}\|_{\mathbf{V}}^2$. Continuity of the bilinear form $(\operatorname{div} \cdot, \cdot)$ is shown in the following lemma.

LEMMA 2.1. *The inequality*

$$|(\operatorname{div} \mathbf{u}, p)| \leq \sqrt{d} \|\mathbf{u}\|_{\mathbf{V}} \|p\|_M$$

holds for all $\mathbf{u} \in \mathbf{V}$, $p \in M$.

Proof. This result immediately follows from the Cauchy inequality and the estimate $\|\nu^{\frac{1}{2}} \operatorname{div} \mathbf{u}\| \leq \sqrt{d} \|\mathbf{u}\|_{\mathbf{V}}$ for $\mathbf{u} \in \mathbf{V}$. \square

We now derive a uniform (w.r.t. ν) infsup property corresponding to the problem (1.6). It generalizes the well-known Nečas inequality:

$$c(\Omega) \|p\| \leq \|\nabla p\|_{-1} := \sup_{\mathbf{u} \in \mathbf{V}} \frac{(\operatorname{div} \mathbf{u}, p)}{\|\nabla \mathbf{u}\|} \quad \forall p \in L^2(\Omega) : (p, 1) = 0, \quad (2.1)$$

with $c(\Omega) > 0$. We will need an equivalent form of (2.1): for any $p \in L^2(\Omega)$ such that $(p, 1) = 0$ there exists $\mathbf{u} \in \mathbf{V}$ such that

$$\|p\|^2 = (\operatorname{div} \mathbf{u}, p) \quad \text{and} \quad c(\Omega) \|\nabla \mathbf{u}\| \leq \|p\|. \quad (2.2)$$

We introduce the piecewise constant function

$$\bar{p} = \begin{cases} |\Omega_1|^{-1} & \text{on } \Omega_1 \\ -\varepsilon |\Omega_2|^{-1} & \text{on } \Omega_2. \end{cases} \quad (2.3)$$

and the one-dimensional subspace $M_0 := \operatorname{span}\{\bar{p}\}$ of M . The proof of theorem 2.2 relies on the $(\cdot, \cdot)_M$ -orthogonal decomposition $M = M_0 \oplus M_0^\perp$. For $p \in M$ we use the notation

$$p = p_0 + p_0^\perp, \quad p_0 \in M_0, \quad p_0^\perp \in M_0^\perp \quad (2.4)$$

One easily verifies that

$$M_0^\perp = \left\{ p \in M \mid \int_{\Omega_1} p \, dx = \int_{\Omega_2} p \, dx = 0 \right\}$$

THEOREM 2.2. *There exists a constant $C > 0$ independent of ν such that*

$$\sup_{\mathbf{u} \in \mathbf{V}} \frac{(\operatorname{div} \mathbf{u}, p)}{\|\mathbf{u}\|_{\mathbf{V}}} \geq C \|p\|_M \quad \text{for all } p \in M$$

Proof. Fix an arbitrary $p \in M$. We first consider the component p_0^\perp from the decomposition $p = p_0 + p_0^\perp$ in (2.4). Since $p_0^\perp|_{\Omega_k} \in L^2(\Omega_k)$ and $(p_0^\perp, 1)_{\Omega_k} = 0$ for $k = 1, 2$, we can apply the Nečas inequality in the form (2.2) in each subdomain. Thus there exists a function $\mathbf{u}_1 \in H_0^1(\Omega_1)^d$ such that the following relations hold with a constant $c(\Omega_1) > 0$:

$$\|p_0^\perp\|_{\Omega_1}^2 = (\operatorname{div} \mathbf{u}_1, p_0^\perp)_{\Omega_1} \quad \text{and} \quad c(\Omega_1) \|\nabla \mathbf{u}_1\|_{\Omega_1} \leq \|p_0^\perp\|_{\Omega_1} \quad (2.5)$$

Similarly, using a scaling argument, it follows that there exists $\mathbf{u}_2 \in H_0^1(\Omega_2)^d$ such that

$$\|\varepsilon^{-\frac{1}{2}} p_0^\perp\|_{\Omega_2}^2 = (\operatorname{div} \mathbf{u}_2, p_0^\perp)_{\Omega_2}, \quad c(\Omega_2) \|\varepsilon^{\frac{1}{2}} \nabla \mathbf{u}_2\|_{\Omega_2} \leq \|\varepsilon^{-\frac{1}{2}} p_0^\perp\|_{\Omega_2}, \quad (2.6)$$

with $c(\Omega_2) > 0$. Extending \mathbf{u}_1 and \mathbf{u}_2 by zero on the whole domain Ω and taking a sum of (2.5) and (2.6) we get

$$\|p_0^\perp\|_M^2 = (\operatorname{div} \tilde{\mathbf{u}}, p_0^\perp) \quad \text{and} \quad c_1 \|\tilde{\mathbf{u}}\|_{\mathbf{V}} \leq \|p_0^\perp\|_M, \quad \tilde{\mathbf{u}} := \mathbf{u}_1 + \mathbf{u}_2 \quad (2.7)$$

with $c_1 = \min\{c(\Omega_1), c(\Omega_2)\}$.

For the component p_0 we define $\tilde{p}_0 := \nu^{-1} p_0$. Note that $(\tilde{p}_0, 1) = (p_0, 1)_M = 0$ and thus we can use the Nečas inequality in Ω . Hence there exists $\tilde{\mathbf{u}} \in H_0^1(\Omega)^d$ such that

$$\|\tilde{p}_0\|^2 = (\operatorname{div} \tilde{\mathbf{u}}, \tilde{p}_0) \quad \text{and} \quad c(\Omega) \|\nabla \tilde{\mathbf{u}}\| \leq \|\tilde{p}_0\|. \quad (2.8)$$

Due to the definition of M_0 we obtain

$$\|p_0\|_M^2 = C(\varepsilon, \Omega) \|\tilde{p}_0\|^2 \quad \text{and} \quad (\operatorname{div} \tilde{\mathbf{u}}, p_0) = C(\varepsilon, \Omega) (\operatorname{div} \tilde{\mathbf{u}}, \tilde{p}_0)$$

with $C(\varepsilon, \Omega) = \frac{\varepsilon|\Omega_1| + |\Omega_2|}{|\Omega_1| + |\Omega_2|}$. Note that

$$C(\varepsilon, \Omega) \geq \tilde{c}(\Omega) \max\{1, \varepsilon\}, \quad \tilde{c}(\Omega) := \min\left\{\frac{|\Omega_1|}{|\Omega|}, \frac{|\Omega_2|}{|\Omega|}\right\} \quad (2.9)$$

From this and (2.8) we deduce

$$\|p_0\|_M^2 = (\operatorname{div} \tilde{\mathbf{u}}, p_0) \quad \text{and} \quad c_3 \max\{1, \sqrt{\varepsilon}\} \|\nabla \tilde{\mathbf{u}}\| \leq \|p_0\|_M, \quad (2.10)$$

with $c_3 = c(\Omega) \tilde{c}(\Omega)^{\frac{1}{2}}$ a constant independent of ν . We also have:

$$(\operatorname{div} \tilde{\mathbf{u}}, p_0) = 0, \quad \|\nu^{\frac{1}{2}} \operatorname{div} \tilde{\mathbf{u}}\| \leq \sqrt{d} \|\tilde{\mathbf{u}}\|_{\mathbf{V}}, \quad \|\tilde{\mathbf{u}}\|_{\mathbf{V}} \leq \max\{1, \sqrt{\varepsilon}\} \|\nabla \tilde{\mathbf{u}}\| \leq c_3^{-1} \|p_0\|_M$$

Using this and the results in (2.7) and (2.10) we get for arbitrary $\alpha > 0$

$$\begin{aligned} (\operatorname{div}(\alpha \tilde{\mathbf{u}} + \tilde{\mathbf{u}}), p) &= \alpha \|p_0^\perp\|_M^2 + \|p_0\|_M^2 + (\operatorname{div} \tilde{\mathbf{u}}, p_0^\perp) \\ &\geq \alpha \|p_0^\perp\|_M^2 + \|p_0\|_M^2 - c_3^{-1} \sqrt{d} \|p_0\|_M \|p_0^\perp\|_M \\ &\geq \frac{1}{2} \|p\|_M^2 \quad \text{if} \quad \alpha \geq \frac{1}{2} \left(1 + \frac{d}{c_3^2}\right) =: \alpha_0 \end{aligned}$$

Thus if we take $\mathbf{u} = \alpha_0 \tilde{\mathbf{u}} + \tilde{\mathbf{u}}$ we get

$$\|p\|_M^2 \leq 2(\operatorname{div} \mathbf{u}, p) \quad \text{and} \quad \|\mathbf{u}\|_{\mathbf{V}}^2 \leq 2(\alpha_0^2 \|\tilde{\mathbf{u}}\|_{\mathbf{V}}^2 + \|\tilde{\mathbf{u}}\|_{\mathbf{V}}^2) \leq c \|p\|_M^2,$$

with a constant c independent of ν . \square

It follows that we have ellipticity of the bilinear form $(\nu \nabla \cdot, \nabla \cdot)$, continuity of the bilinear forms $(\nu \nabla \cdot, \nabla \cdot)$ and $(\operatorname{div} \cdot, \cdot)$, and the infsup property in the norms $\|\cdot\|_{\mathbf{V}}$ and $\|\cdot\|_M$ with constants that are independent of ν . Thus we have uniform (w.r.t. ν) well-posedness of the continuous variational Stokes problem (1.6) in these norms. Using standard arguments (cf. [17, 10]) it can be shown that the problem (1.6) has a unique solution and that the a priori estimate

$$(\|u\|_{\mathbf{V}}^2 + \|p\|_M^2)^{\frac{1}{2}} \leq c \|\mathbf{f}\|_{\mathbf{V}} \quad (2.11)$$

holds with a constant c independent of \mathbf{f} and of ν .

REMARK 1. We comment on the dependence of the dual norm $\|\mathbf{f}\|_{\mathbf{V}'}$ in (2.11) on the viscosity parameter ν . For this we need the Poincare type inequality

$$\|\nu^{\frac{1}{2}}\mathbf{v}\| \leq C_P\|\mathbf{v}\|_{\mathbf{V}}, \quad \text{for all } \mathbf{v} \in \mathbf{V}. \quad (2.12)$$

The optimal constant C_P in (2.12) is uniformly bounded w.r.t. ν if one of the following conditions is satisfied:

$$\text{meas}(\partial\Omega_k \cap \partial\Omega) > 0 \quad \text{for } k = 1, 2 \quad (2.13)$$

$$\text{meas}(\partial\Omega_1 \cap \partial\Omega) > 0 \quad \text{and } \varepsilon \leq C \quad (2.14)$$

The fact that the condition (2.13) is sufficient for uniform boundedness of C_P follows from lemma 1 and (the proof of) lemma 7 in [22]. The fact that the condition (2.14) is sufficient for uniform boundedness of C_P is proved with the following argument. Due to (2.14) \mathbf{u} vanishes on a part of $\partial\Omega_1$ with nonzero measure, hence

$$\|\mathbf{u}\|_{\Omega_1} \leq c\|\nabla\mathbf{u}\|_{\Omega_1}. \quad (2.15)$$

holds. Therefore

$$\|\mathbf{u}|_{\Gamma}\| = \|\mathbf{u}|_{\partial\Omega_1}\| \leq c\|\nabla\mathbf{u}\|_{\Omega_1}. \quad (2.16)$$

In the subdomain Ω_2 we have

$$\varepsilon\|\mathbf{u}\|_{\Omega_2}^2 \leq c\varepsilon(\|\nabla\mathbf{u}\|_{\Omega_2}^2 + \|\mathbf{u}|_{\partial\Omega_2}\|^2) = c\varepsilon(\|\nabla\mathbf{u}\|_{\Omega_2}^2 + \|\mathbf{u}|_{\Gamma}\|^2). \quad (2.17)$$

Inequalities (2.15), (2.16), and (2.17) give the inequality (2.12) with a constant C independent of ε and hence independent of ν .

Assume now that $\mathbf{f} \in L^2(\Omega)^d$ and that one of conditions (2.13),(2.14) holds. Then the Cauchy inequality and (2.12) immediately yield the a-priori estimate

$$(\|\mathbf{u}\|_{\mathbf{V}}^2 + \|p\|_M^2)^{\frac{1}{2}} \leq cC_P\|\nu^{-\frac{1}{2}}\mathbf{f}\|, \quad (2.18)$$

with cC_P independent of \mathbf{f} and of ν .

3. Finite element discretization. In this section we consider the discretization of the variational Stokes problem using a family of pairs of conforming finite element spaces. For this we assume a family of triangulations $\{\mathcal{T}_h\}$ in the sense of [11, 12]. An important assumption for our analysis is that each triangulation \mathcal{T}_h is conforming w.r.t. the two subdomains Ω_1, Ω_2 in the following sense:

$$\exists \mathcal{T}_h^{(i)} \subset \mathcal{T}_h : \quad \cup\{T \mid T \in \mathcal{T}_h^{(i)}\} = \bar{\Omega}_i, \quad i = 1, 2 \quad (3.1)$$

This assumption is easily fulfilled if Ω_1 and Ω_2 are polyhedral subdomains.

REMARK 2. In computational fluid dynamics for two-phase flow problems it is (more) realistic to assume that $\Gamma = \partial\Omega_1 \cap \partial\Omega_2$ is smooth. Then the assumption (3.1) in general does not hold. However, in such applications it is common practice to approximate Γ by a polyhedral discrete interface Γ_h . In such a setting the assumption (3.1) may still make sense. As far as we know no rigorous analysis is available which for the (Navier)-Stokes equations shows the effect of approximating the smooth interface Γ by a piecewise smooth interface Γ_h . A theoretical analysis of this effect for a Poisson interface problem can be found in [15]. The results in [15], however, are not robust with respect to the jump in the diffusion coefficient.

We assume a pair of finite element spaces $\mathbf{V}_h \subset \mathbf{V}$ and $Q_h \subset L_0^2(\Omega) = \{p \in L^2(\Omega) \mid (p, 1) = 0\}$ that is LBB stable with a constant $\hat{\beta}$ independent of h :

$$\inf_{q_h \in Q_h} \sup_{\mathbf{v}_h \in \mathbf{V}_h} \frac{(\operatorname{div} \mathbf{v}_h, q_h)}{\|\nabla \mathbf{v}_h\| \|q_h\|} \geq \hat{\beta} > 0. \quad (3.2)$$

Note that due to the different normalization in the space M (namely $(p, 1)_M = 0$) we in general have $Q_h \not\subset M$. To maintain conformity we use the space

$$M_h = \{p_h = \tilde{p}_h + \alpha 1 \mid \tilde{p}_h \in Q_h, \quad \alpha \in \mathbb{R} \text{ such that } (p_h, 1)_M = 0\}.$$

Note that $M_h \subset M$ and that functions in M_h and in Q_h only differ by a constant.

For the analysis in this section it is convenient (but not necessary) to introduce the bilinear form $a : (\mathbf{V} \times M) \times (\mathbf{V} \times M) \rightarrow \mathbb{R}$

$$a(\mathbf{u}, p; \mathbf{v}, q) := (\nu \nabla \mathbf{u}, \nabla \mathbf{v}) - (\operatorname{div} \mathbf{v}, p) + (\operatorname{div} \mathbf{u}, q) \quad (3.3)$$

and formulate the discrete problem as follows: find $\{\mathbf{u}_h, p_h\} \in \mathbf{V}_h \times M_h$ such that

$$a(\mathbf{u}_h, p_h; \mathbf{v}_h, q_h) = (\mathbf{f}, \mathbf{v}_h) \quad \text{for all } \{\mathbf{v}_h, q_h\} \in \mathbf{V}_h \times M_h. \quad (3.4)$$

In this section we will analyze continuity (theorem 3.4) and discrete stability (theorem 3.5) of the bilinear form $a(\cdot, \cdot)$. The estimates are uniform with respect to the mesh size parameter h and the diffusion coefficient ν . As a corollary we then obtain a uniform discretization error bound.

In the proof of the discrete infsup condition below we will use a decomposition which is similar, but not identical to the one from the previous section. Let $\bar{p}_h \in M_h$ be the M -orthogonal projection of \bar{p} on M_h ,

$$(\bar{p} - \bar{p}_h, q_h)_M = 0 \quad \text{for all } q_h \in M_h$$

and define the one-dimensional subspace $M_{0,h} := \operatorname{span}(\bar{p}_h)$ of M_h . This induces an $(\cdot, \cdot)_M$ -orthogonal decomposition of M (and also of M_h): $M = M_{0,h} \oplus M_{0,h}^\perp$, and for $p \in M$ we use the notation

$$p = p_{0,h} + p_{0,h}^\perp, \quad p_{0,h} \in M_{0,h}, \quad p_{0,h}^\perp \in M_{0,h}^\perp. \quad (3.5)$$

We will need the following elementary result

LEMMA 3.1. *For all $p_h = p_{0,h} + p_{0,h}^\perp \in M_h$ we have*

$$(p_{0,h}^\perp, 1)_{\Omega_k} = 0 \quad \text{for } k = 1, 2$$

Proof. First note that by definition $(p_{0,h}^\perp, \bar{p}_h)_M = 0$. Using $(p_{0,h}^\perp, \bar{p}_h)_M = (p_{0,h}^\perp, \bar{p})_M$ we then get

$$0 = \frac{1}{|\Omega_1|} (p_{0,h}^\perp, 1)_{\Omega_1} - \frac{1}{|\Omega_2|} (p_{0,h}^\perp, 1)_{\Omega_2} \quad (3.6)$$

Since $p_{0,h}^\perp \in M$ we have $(p_{0,h}^\perp, 1)_M = 0$ and thus

$$(p_{0,h}^\perp, 1)_{\Omega_1} + \varepsilon^{-1} (p_{0,h}^\perp, 1)_{\Omega_2} = 0 \quad (3.7)$$

Combination of (3.6) and (3.7) proves the lemma. \square

In the analysis below we use the quantity

$$\mu_h := \frac{\|\bar{p} - \bar{p}_h\|_M}{\|\bar{p}\|_M} = \inf_{q_h \in M_h} \frac{\|\bar{p} - q_h\|_M}{\|\bar{p}\|_M} \quad (3.8)$$

which measures the error made by approximating \bar{p} in the finite element pressure space. We will need that this quantity tends to zero (with a rate independent of ν) if h tends to zero. This issue is discussed in the following remark.

REMARK 3. We always have $\mu_h \leq 1$. Furthermore, $\mu_h = 0$ if M_h contains piecewise constant finite elements. We now consider the case where M_h consists of continuous functions which are piecewise polynomial on the triangulation \mathcal{T}_h . Let Γ_h be the set of elements from \mathcal{T}_h which have at least one vertex lying on the interface Γ . We assume $|\Gamma_h| \leq c_T h$ with a constant c_T independent of h . Let $I(\bar{p})$ be the continuous piecewise linear function which has value $\bar{p}|_{\Omega_1}$ at all vertices in $\bar{\Omega}_1 \setminus \bar{\Gamma}$, value $\bar{p}|_{\Omega_2}$ at all vertices in $\bar{\Omega}_2 \setminus \bar{\Gamma}$ and $I(\bar{p}) = 0$ on $\bar{\Gamma}$. Note that $I(\bar{p}) \neq \bar{p}$ only in Γ_h . A simple computation yields

$$\begin{aligned} \|\bar{p}\|_M^2 &= |\Omega_1|^{-1} + \varepsilon |\Omega_2|^{-1} \\ \|\bar{p} - I(\bar{p})\|_M^2 &= \int_{\Gamma_h} \nu^{-1} (\bar{p} - I(\bar{p}))^2 dx \leq |\Gamma_h| (|\Omega_1|^{-2} + \varepsilon |\Omega_2|^{-2}) \end{aligned}$$

Thus we obtain

$$\frac{\|\bar{p} - I(\bar{p})\|_M}{\|\bar{p}\|_M} \leq c_T^{\frac{1}{2}} (|\Omega_1|^{-1} + |\Omega_2|^{-1})^{\frac{1}{2}} h^{\frac{1}{2}} \quad (3.9)$$

Note that $I(\bar{p})$ does not necessarily satisfy the orthogonality condition $(I(\bar{p}), 1)_M = 0$. Therefore we introduce the continuous piecewise linear function $q_h := I(\bar{p}) - \alpha 1$ with $\alpha = \frac{(I(\bar{p}), 1)_M}{\|1\|_M^2}$, i.e., $(q_h, 1)_M = 0$ and thus $q_h \in M_h$. Note that

$$|\alpha| = \frac{|(I(\bar{p}) - \bar{p}, 1)_M|}{\|1\|_M^2} \leq \frac{\|I(\bar{p}) - \bar{p}\|_M}{\|1\|_M}$$

We then finally get

$$\begin{aligned} \mu_h &\leq \frac{\|\bar{p} - q_h\|_M}{\|\bar{p}\|_M} \leq \frac{\|\bar{p} - I(\bar{p})\|_M}{\|\bar{p}\|_M} + \frac{|\alpha| \|1\|_M}{\|\bar{p}\|_M} \\ &\leq 2 \frac{\|\bar{p} - I(\bar{p})\|_M}{\|\bar{p}\|_M} \leq 2c_T^{\frac{1}{2}} (|\Omega_1|^{-1} + |\Omega_2|^{-1})^{\frac{1}{2}} h^{\frac{1}{2}} \end{aligned}$$

and thus we conclude that $\mu_h \leq ch^{\frac{1}{2}}$ holds with a constant c independent of ν .

For the analysis of the discrete infsup property we need the following result:

LEMMA 3.2. *For every $p_{0,h} \in M_{0,h}$ there exist $p_0 \in M_0$ such that*

$$\|p_{0,h} - p_0\|_M = \mu_h \|p_0\|_M \quad (3.10)$$

$$\|p_{0,h}\|_M = \sqrt{1 - \mu_h^2} \|p_0\|_M \quad (3.11)$$

Proof. For $p_{0,h} \in M_{0,h}$ we have $p_{0,h} = \alpha \bar{p}_h$ with $\alpha \in \mathbb{R}$. We set $p_0 = \alpha \bar{p}$. Since \bar{p}_h is the M -orthogonal projection of \bar{p} on $M_{0,h}$, $p_{0,h}$ is the M -orthogonal projection of p_0 on $M_{0,h}$. This choice of p_0 implies (3.10) by definition of μ_h . The result in (3.11) follows from

$$\|p_0\|_M^2 = \|p_{0,h} - p_0\|_M^2 + \|p_{0,h}\|_M^2 = \mu_h^2 \|p_0\|_M^2 + \|p_{0,h}\|_M^2$$

□

We need an additional assumption on \mathbf{V}_h . Consider $\tilde{p} = \nu^{-1} \bar{p}$ with \bar{p} defined in (2.3). \tilde{p} has zero mean: $(\tilde{p}, 1) = 0$. We assume that \mathbf{V}_h is such that there is a constant $\hat{\beta}_c > 0$ independent of h such that

$$\sup_{\mathbf{v}_h \in \mathbf{V}_h} \frac{(\operatorname{div} \mathbf{v}_h, \tilde{p})}{\|\nabla \mathbf{v}_h\|} \geq \hat{\beta}_c \|\tilde{p}\|. \quad (3.12)$$

REMARK 4. Assumption (3.12) is rather weak. We briefly discuss two cases in which this assumption is satisfied. Let h_0 be the mesh size parameter corresponding to the coarsest triangulation. Then (3.12) trivially holds for \mathbf{V}_{h_0} with a constant $\hat{\beta}_c = \hat{\beta}_c(h_0)$. If the family of spaces $\{\mathbf{V}_h\}_{h \leq h_0}$ is *nested* then (3.12) with $\hat{\beta}_c = \hat{\beta}_c(h_0)$ holds for any \mathbf{V}_h . The second case is when M_h contains piecewise constant elements. Then (3.12) immediately follows from (3.2).

We now prove a discrete infsup stability result, which is a main result of this paper:

THEOREM 3.3. *There exist constants $C_1 > 0, C_2 > 0$ independent of ν and h such that*

$$\text{if } \mu_h \leq C_1 \text{ then} \quad (3.13)$$

$$\sup_{\mathbf{u}_h \in \mathbf{V}_h} \frac{(\operatorname{div} \mathbf{u}_h, p_h)}{\|\mathbf{u}_h\|_{\mathbf{V}}} \geq C_2 \|p_h\|_M \quad \text{for all } p_h \in M_h \quad (3.14)$$

Proof. The proof is based on similar arguments as used in the proof of theorem 2.2. All the constants that appear in the proof are independent of ν and of h .

Take an arbitrary $p_h \in M_h$. We first consider the component $p_{0,h}^\perp$ from the decomposition $p_h = p_{0,h} + p_{0,h}^\perp$ in (3.5). From lemma 3.1 it follows that $(p_{0,h}^\perp, 1)_{\Omega_k} = 0$ for $k = 1, 2$ and thus we can use the LBB property (3.2) in each subdomain. Hence there exists a function $\mathbf{u}_1 \in \mathbf{V}_h$ with $\mathbf{u}_1 = 0$ on Ω_2 such that the following relations hold with a constant $c_1 > 0$:

$$\|p_{0,h}^\perp\|_{\Omega_1}^2 = (\operatorname{div} \mathbf{u}_1, p_{0,h}^\perp)_{\Omega_1} \quad \text{and} \quad c_1 \|\nabla \mathbf{u}_1\|_{\Omega_1} \leq \|p_{0,h}^\perp\|_{\Omega_1}. \quad (3.15)$$

Similarly, using a scaling argument, it follows that there exists $\mathbf{u}_2 \in \mathbf{V}_h$ with $\mathbf{u}_2 = 0$ on Ω_1 such that

$$\|\varepsilon^{-\frac{1}{2}} p_{0,h}^\perp\|_{\Omega_2}^2 = (\operatorname{div} \mathbf{u}_2, p_{0,h}^\perp)_{\Omega_2} \quad \text{and} \quad c_2 \|\varepsilon^{\frac{1}{2}} \nabla \mathbf{u}_2\|_{\Omega_2} \leq \|\varepsilon^{-\frac{1}{2}} p_{0,h}^\perp\|_{\Omega_2} \quad (3.16)$$

with a constant $c_2 > 0$. Taking a sum of (3.15) and (3.16) we get for $\tilde{\mathbf{u}}_h := \mathbf{u}_1 + \mathbf{u}_2$:

$$\|p_{0,h}^\perp\|_M^2 = (\operatorname{div} \tilde{\mathbf{u}}_h, p_{0,h}^\perp) \quad \text{and} \quad \hat{c} \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}} \leq \|p_{0,h}^\perp\|_M, \quad (3.17)$$

with $\hat{c} := \min\{c_1, c_2\}$. We now consider the component $p_{0,h}$. Take $p_0 \in M_0$ as in lemma 3.2. Then we have

$$\|p_{0,h} - p_0\|_M = \mu_h \|p_0\|_M, \quad \|p_{0,h}\|_M = \sqrt{1 - \mu_h^2} \|p_0\|_M \quad (3.18)$$

For $\tilde{p}_0 := \nu^{-1}p_0$ we get $(\tilde{p}_0, 1) = (p_0, 1)_M = 0$. From assumption (3.12) it follows that there exists $\tilde{\mathbf{u}}_h \in \mathbf{V}_h$ such that

$$\|\tilde{p}_0\|^2 = (\operatorname{div} \tilde{\mathbf{u}}_h, \tilde{p}_0) \quad \text{and} \quad \hat{\beta}_c \|\nabla \tilde{\mathbf{u}}_h\| \leq \|\tilde{p}_0\|$$

Using the same arguments as in the proof of theorem 2.2 one can show that

$$\|p_0\|_M^2 = (\operatorname{div} \tilde{\mathbf{u}}_h, p_0) \quad \text{and} \quad \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}} \leq c_3^{-1} \|p_0\|_M \quad (3.19)$$

with $c_3 = \hat{\beta}_c \tilde{c}(\Omega)^{\frac{1}{2}}$ and $\tilde{c}(\Omega)$ the constant from (2.9). Also note that $(\operatorname{div} \tilde{\mathbf{u}}_h, p_0) = 0$. Using this and the results in (3.17), (3.19) we obtain for arbitrary $\alpha > 0$:

$$\begin{aligned} (\operatorname{div}(\alpha \tilde{\mathbf{u}}_h + \tilde{\mathbf{u}}_h), p_h) &= \alpha (\operatorname{div} \tilde{\mathbf{u}}_h, p_0 + (p_{0,h} - p_0) + p_{0,h}^\perp) \\ &\quad + (\operatorname{div} \tilde{\mathbf{u}}_h, p_0 + (p_{0,h} - p_0) + p_{0,h}^\perp) \\ &= \alpha \|p_{0,h}^\perp\|_M^2 + \alpha (\operatorname{div} \tilde{\mathbf{u}}_h, p_{0,h} - p_0) \\ &\quad + \|p_0\|_M^2 + (\operatorname{div} \tilde{\mathbf{u}}_h, (p_{0,h} - p_0) + p_{0,h}^\perp) \end{aligned}$$

We assume that $\mu_h^2 \leq \frac{1}{2}$. From (3.18) we then get

$$\|p_{0,h}\|_M^2 \leq \|p_0\|_M^2 \leq 2 \|p_{0,h}\|_M^2$$

and using $\|\nu^{\frac{1}{2}} \operatorname{div} \mathbf{v}_h\| \leq \sqrt{d} \|\mathbf{v}_h\|_{\mathbf{V}}$ for $\mathbf{v}_h \in \mathbf{V}_h$ we obtain

$$\begin{aligned} (\operatorname{div}(\alpha \tilde{\mathbf{u}}_h + \tilde{\mathbf{u}}_h), p_h) &\geq \alpha \|p_{0,h}^\perp\|_M^2 + \|p_{0,h}\|_M^2 - \alpha \sqrt{d} \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}} \|p_{0,h} - p_0\|_M \\ &\quad - \sqrt{d} \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}} (\|p_{0,h} - p_0\|_M + \|p_{0,h}^\perp\|_M) \\ &\geq \alpha \|p_{0,h}^\perp\|_M^2 + \|p_{0,h}\|_M^2 - \alpha \sqrt{2d} \hat{c}^{-1} \mu_h \|p_{0,h}^\perp\|_M \|p_{0,h}\|_M \\ &\quad - c_3^{-1} \sqrt{2d} \|p_{0,h}\|_M (\sqrt{2} \mu_h \|p_{0,h}\|_M + \|p_{0,h}^\perp\|_M) \\ &= \alpha \|p_{0,h}^\perp\|_M^2 + (1 - 2c_3^{-1} \sqrt{d} \mu_h) \|p_{0,h}\|_M^2 \\ &\quad - \sqrt{2d} (\alpha \hat{c}^{-1} \mu_h + c_3^{-1}) \|p_{0,h}\|_M \|p_{0,h}^\perp\|_M \end{aligned}$$

We take $\mu_h \leq \frac{c_3}{4\sqrt{d}}$. Then $1 - 2c_3^{-1} \sqrt{d} \mu_h \geq \frac{1}{2}$ holds and we obtain, using the Cauchy inequality,

$$(\operatorname{div}(\alpha \tilde{\mathbf{u}}_h + \tilde{\mathbf{u}}_h), p_h) \geq (\alpha - 2d(\alpha \hat{c}^{-1} \mu_h + c_3^{-1})^2) \|p_{0,h}^\perp\|_M^2 + \frac{1}{4} \|p_{0,h}\|_M^2$$

We now take

$$\mu_h \leq \frac{1}{\alpha_0}, \quad \alpha_0 := \frac{1}{4} + 2d(\hat{c}^{-1} + c_3^{-1})^2$$

and then for $\mathbf{u}_h := \alpha_0 \tilde{\mathbf{u}}_h + \tilde{\mathbf{u}}_h$ we obtain

$$\begin{aligned} (\operatorname{div} \mathbf{u}_h, p_h) &\geq \frac{1}{4} \|p_h\|_M^2 \\ \|\mathbf{u}_h\|_{\mathbf{V}}^2 &\leq 2(\alpha_0^2 \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}}^2 + \|\tilde{\mathbf{u}}_h\|_{\mathbf{V}}^2) \leq C \|p_h\|_M^2 \end{aligned}$$

with a constant C independent of h and ν .

Hence for $C_1 = \min \left\{ \frac{1}{\sqrt{2}}, \frac{c_3}{4\sqrt{d}}, \frac{1}{\alpha_0} \right\}$ we have the desired result. \square

Note that for h sufficiently small (independent of ν) the condition $\mu_h \leq C_1$ in (3.13) is fulfilled, cf. remark 3.

We now use standard arguments to derive continuity and stability results for the bilinear form $a(\cdot, \cdot)$. For completeness we also present the proofs. We introduce the product norm

$$\| \mathbf{u}, p \| = \left(\| \mathbf{u} \|_{\mathbf{V}}^2 + \| p \|_M^2 \right)^{\frac{1}{2}} \quad \{ \mathbf{u}, p \} \in \mathbf{V} \times M$$

From $(\nu \nabla \mathbf{u}, \nabla \mathbf{u}) = \| \mathbf{u} \|_{\mathbf{V}}^2$ and the result in lemma 2.1 we immediately obtain the following continuity result:

THEOREM 3.4. *There exists a constant C independent of ν such that*

$$a(\mathbf{u}, p; \mathbf{v}, q) \leq C \| \mathbf{u}, p \| \| \mathbf{v}, q \|$$

for all $\{ \mathbf{u}, p \}, \{ \mathbf{v}, q \} \in \mathbf{V} \times M$.

A discrete infsup result is presented in the next theorem:

THEOREM 3.5. *Assume that the condition (3.13) is satisfied. There exists a constant $c > 0$ independent of h and of ν such that*

$$\sup_{\{ \mathbf{v}_h, q_h \} \in \mathbf{V}_h \times M_h} \frac{a(\mathbf{u}_h, p_h; \mathbf{v}_h, q_h)}{\| \mathbf{v}_h, q_h \|} \geq c \| \mathbf{u}_h, p_h \| \quad \forall \{ \mathbf{u}_h, p_h \} \in \mathbf{V}_h \times M_h$$

Proof. Take $\{ \mathbf{u}_h, p_h \} \in \mathbf{V}_h \times M_h$. From the infsup result in theorem 3.3 it follows that there exists $\mathbf{z}_h \in \mathbf{V}_h$ such that $\| \mathbf{z}_h \|_{\mathbf{V}} = \| p_h \|_M$ and $-(\operatorname{div} \mathbf{z}_h, p_h) \geq c \| p_h \|_M^2$ with $c > 0$. Now take $\mathbf{v}_h := \mathbf{u}_h + c \mathbf{z}_h$, $q_h := p_h$. We then get

$$\begin{aligned} a(\mathbf{u}_h, p_h; \mathbf{u}_h, p_h) &= \| \mathbf{u}_h \|_{\mathbf{V}}^2, \\ a(\mathbf{u}_h, p_h; \mathbf{z}_h, 0) &\geq \frac{c}{2} \| p_h \|_M^2 - \frac{1}{2c} \| \mathbf{u}_h \|_{\mathbf{V}}^2. \end{aligned}$$

We multiply the second inequality by c and add it to the first one. This gives

$$a(\mathbf{u}_h, p_h; \mathbf{u}_h + c \mathbf{z}_h, p_h) \geq \frac{1}{2} \| \mathbf{u}_h \|_{\mathbf{V}}^2 + \frac{c^2}{2} \| p_h \|_M^2 \geq c_1 \| \mathbf{u}_h, p_h \|^2, \quad (3.20)$$

with $c_1 = \frac{1}{2} \min \{ 1, c^2 \}$. Now note

$$\begin{aligned} \| \mathbf{v}_h, p_h \|^2 &\leq 2(\| \mathbf{u}_h \|_{\mathbf{V}}^2 + c^2 \| \mathbf{z}_h \|_{\mathbf{V}}^2) + \| p_h \|_M^2 \\ &= 2 \| \mathbf{u}_h \|_{\mathbf{V}}^2 + (2c^2 + 1) \| p_h \|_M^2 \leq 2(c^2 + 1) \| \mathbf{u}_h, p_h \|^2 \end{aligned} \quad (3.21)$$

Combination of (3.20) and (3.21) completes the proof. \square

As for the continuous problem we get as a direct corollary that the discrete problem (3.4) has a unique solution $\{ \mathbf{u}_h, p_h \}$ and the inequality

$$\| \mathbf{u}_h, p_h \| \leq c^{-1} \| \mathbf{f} \|_{\mathbf{V}'_h}$$

holds, with the constant c from theorem 3.5. Moreover, if $\mathbf{f} \in L^2(\Omega)^d$, then using the Cauchy inequality and the Poincare inequality (2.12) we obtain the a-priori estimate:

$$\| \mathbf{u}_h, p_h \| \leq c^{-1} C_P \| \nu^{-\frac{1}{2}} \mathbf{f} \|. \quad (3.22)$$

We refer to remark 1 for a discussion of the dependence of the Poincare “constant” on ν .

Using the continuity result in theorem 3.4 and the infsup result in theorem 3.5 we can prove a discretization error bound using standard arguments.

THEOREM 3.6. *Let $\{\mathbf{u}, p\}$ be the solution of the continuous problem (1.6) and $\{\mathbf{u}_h, p_h\}$ be the solution of the discrete problem (3.4). Assume that the condition (3.13) is satisfied. There exists a constant C independent of h and of ν such that the following holds:*

$$\|\|\mathbf{u} - \mathbf{u}_h, p - p_h\|\| \leq C \min_{\mathbf{v}_h \in \mathbf{V}_h} \min_{q_h \in M_h} \|\|\mathbf{u} - \mathbf{v}_h, p - q_h\|\|. \quad (3.23)$$

Proof. For arbitrary $\mathbf{v}_h \in \mathbf{V}_h, q_h \in M_h$ define $\mathbf{e} := \mathbf{u} - \mathbf{v}_h, \mathbf{e}_h = \mathbf{u}_h - \mathbf{v}_h, g := p - q_h, g_h := p_h - q_h$. The Galerkin orthogonality property yields

$$a(\mathbf{e}_h, g_h; \mathbf{z}_h, r_h) = a(\mathbf{e}, g; \mathbf{z}_h, r_h) \quad \text{for all } \{\mathbf{z}_h, r_h\} \in \mathbf{V}_h \times M_h.$$

Using this in combination with the continuity and infsup results we obtain, for suitable $\{\mathbf{z}_h, r_h\} \in \mathbf{V}_h \times M_h$:

$$\begin{aligned} \|\|\mathbf{e}_h, g_h\|\| &\leq c^{-1} \frac{a(\mathbf{e}_h, g_h; \mathbf{z}_h, r_h)}{\|\|\mathbf{z}_h, r_h\|\|} = c^{-1} \frac{a(\mathbf{e}, g; \mathbf{z}_h, r_h)}{\|\|\mathbf{z}_h, r_h\|\|} \\ &\leq c^{-1} C \|\|\mathbf{e}, g\|\|. \end{aligned}$$

Now combine this with the triangle inequality $\|\|\mathbf{u} - \mathbf{u}_h, p - p_h\|\| \leq \|\|\mathbf{e}_h, g_h\|\| + \|\|\mathbf{e}, g\|\|$. \square

Based on the result in theorem 3.6 and using approximation properties of the finite element spaces one can derive further bounds for the discretization error. For such an analysis one needs regularity results for the continuous Stokes interface problem. As far as we know, this regularity issue is largely unsolved.

4. Preconditioner for the Schur complement. In this section we analyze convergence properties of iterative solvers for the discretized problem. For this we first introduce the matrix-vector formulation of the discrete problem.

In practice the discrete space M_h for the pressure is constructed by taking a standard finite element space, which we denote by M_h^+ (for example, continuous piecewise linear functions), and then adding an orthogonality condition:

$$M_h = \{p_h \in M_h^+ \mid (p_h, 1)_M = 0\}$$

Note that $\dim(M_h) = \dim(M_h^+) - 1$. Let $n := \dim(\mathbf{V}_h), m := \dim(M_h^+)$. We assume standard (nodal) bases in \mathbf{V}_h and M_h^+ and corresponding isomorphisms

$$J_V : \mathbb{R}^n \rightarrow \mathbf{V}_h, \quad J_M : \mathbb{R}^m \rightarrow M_h^+.$$

Let the stiffness matrices $\mathbf{A} \in \mathbb{R}^{n \times n}, \mathbf{B} \in \mathbb{R}^{m \times n}$ and the mass matrix $\hat{M}_\nu \in \mathbb{R}^{m \times m}$ be given by

$$\begin{aligned} \langle \mathbf{A}x, y \rangle &= (\nu \nabla J_V x, \nabla J_V y) \quad \text{for all } x, y \in \mathbb{R}^n, \\ \langle \mathbf{B}x, y \rangle &= (\text{div } J_V x, J_M y) \quad \text{for all } x \in \mathbb{R}^n, y \in \mathbb{R}^m, \\ \langle \hat{M}_\nu x, y \rangle &= (J_M x, J_M y)_M \quad \text{for all } x, y \in \mathbb{R}^m. \end{aligned} \quad (4.1)$$

Here $\langle \cdot, \cdot \rangle$ denotes the standard Euclidean scalar product. We emphasize that the matrix \hat{M}_ν is the mass matrix with respect to the (weighted L^2) scalar product $(\cdot, \cdot)_M$ and thus may differ very much from the usual mass matrix with respect to the L^2 scalar product (\cdot, \cdot) . After finite element discretization we have a linear system of the form

$$\begin{pmatrix} A & B^T \\ B & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f \\ 0 \end{pmatrix} \quad (4.2)$$

with f such that $\langle f, y \rangle = (f, J_V y)$ for all $y \in \mathbb{R}^n$. The Schur complement is denoted by $S := BA^{-1}B^T$. Note that both S and the matrix in (4.2) are singular and have a one-dimensional kernel. Define the constant vector $e := J_M^{-1}1 = (1, \dots, 1)^T \in \mathbb{R}^m$. Then we have $\ker(S) = \text{span}\{e\}$. Note that

$$(J_{MY}, 1)_M = 0 \Leftrightarrow (J_{MY}, J_M e)_M = 0 \Leftrightarrow \langle \hat{M}_\nu y, e \rangle = 0 \Leftrightarrow \langle y, \hat{M}_\nu e \rangle = 0 \quad (4.3)$$

Hence, with

$$(\hat{M}_\nu e)^\perp := \{y \in \mathbb{R}^m \mid \langle y, \hat{M}_\nu e \rangle = 0\} \quad (4.4)$$

we have $M_h = \{J_{MY} \mid y \in (\hat{M}_\nu e)^\perp\}$ and we get the following matrix-vector representation of the discrete problem (3.4):

$$\text{Find } x \in \mathbb{R}^n, y \in (\hat{M}_\nu e)^\perp \text{ such that (4.2) holds} \quad (4.5)$$

In preconditioned MINRES or (inexact) Uzawa type of iterative solvers for solving this problem one needs preconditioners Q_A of A and Q_S of S . It is known that if for Q_A we take a symmetric multigrid V -cycle then we have (cf. [8, 9, 28])

$$(1 - \sigma_A)Q_A \leq A \leq Q_A,$$

with a constant $\sigma_A < 1$ independent of h and of ν .

Below we show that the mass matrix \hat{M}_ν is an appropriate preconditioner for S . From lemma 2.1 and theorem 3.3 we obtain

$$C_2 \|p_h\|_M \leq \sup_{u_h \in \mathbf{V}_h} \frac{(\text{div } \mathbf{u}_h, p_h)}{\|\mathbf{u}_h\|_{\mathbf{V}}} \leq \sqrt{d} \|p_h\|_M \quad \text{for } p_h \in M_h \quad (4.6)$$

with $C_2 > 0$ independent of h and of ν , provided the condition $\mu_h \leq C_1$ in (3.13) is fulfilled. From the definition of the Schur complement it follows that for arbitrary $y \in \mathbb{R}^m$ we have

$$\langle Sy, y \rangle = \sup_{\mathbf{u}_h \in \mathbf{V}_h} \frac{(\text{div } \mathbf{u}_h, J_{MY})^2}{\|\mathbf{u}_h\|_{\mathbf{V}}^2} \quad (4.7)$$

As a direct consequence of (4.7) and (4.3) we get:

THEOREM 4.1. *Assume that $\mu_h \leq C_1$ in (3.13) holds. For all $y \in (\hat{M}_\nu e)^\perp$ we have*

$$C_2^2 \langle \hat{M}_\nu y, y \rangle \leq \langle Sy, y \rangle \leq d \langle \hat{M}_\nu y, y \rangle$$

with constant C_2 from (4.6).

This theorem shows that the matrix $\hat{M}_\nu^{-1}S$ has a *uniformly bounded spectral condition number on the subspace* $(\bar{M}_\nu e)^\perp$.

One further relevant issue is how to compute of $\hat{M}_\nu^{-1}y$ efficiently. The next lemma shows that either the matrix \bar{M}_ν can be replaced by a cheap diagonal preconditioner or a good approximation of $\hat{M}_\nu^{-1}y$ can be obtained efficiently by applying a preconditioned CG method with a diagonal matrix as preconditioner .

LEMMA 4.2. *Define the diagonal matrix \bar{M}_ν by $(\bar{M}_\nu)_{ii} = \sum_{j=1}^m (\hat{M}_\nu)_{ij}$ (diagonal lumping). Then for all $y \in \mathbb{R}^m$ we have*

$$C_3 \langle \bar{M}_\nu y, y \rangle \leq \langle \hat{M}_\nu y, y \rangle \leq C_4 \langle \bar{M}_\nu y, y \rangle$$

with constants $C_3 > 0$ and C_4 independent of ν and h .

Proof. To show the result it is sufficient to estimate the eigenvalues of $(\bar{M}_\nu|_\tau)^{-1} \hat{M}_\nu|_\tau$ with the local mass matrices $\bar{M}_\nu|_\tau$ and $\hat{M}_\nu|_\tau$ on each element τ of triangulation. On every element ν is constant and thus we can use the result from [27], which yields ν and h - independent bounds on each element. \square

A further elementary observation is

$$\bar{M}_\nu e = \hat{M}_\nu e \tag{4.8}$$

We briefly discuss two known iterative methods for solving the linear system in (4.2). For these methods we will present numerical results in section 5.

A basic method for saddle point problems is the Uzawa method. Applying a block Gaussian elimination step to the system (4.2) yields the equivalent system

$$\begin{pmatrix} I & A^{-1}B^T \\ 0 & S \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} A^{-1}f \\ BA^{-1}f \end{pmatrix}$$

This system can be solved by block backward substitution, which yields the *Uzawa method*:

$$1. \text{ Solve } Az = f \tag{4.9}$$

$$2. \text{ Solve } Sy = Bz, \quad y \in (\hat{M}_\nu e)^\perp \tag{4.10}$$

$$3. \text{ Solve } Ax = z - B^T y \tag{4.11}$$

For the systems in steps 1. and 3. we apply a standard multigrid solver. For the system $Sy = Bz$ we apply a preconditioned CG method (PCG). In each matrix-vector multiplication with S we solve the linear system with A using the multigrid method. For the iterands y^1, y^2, \dots , that are computed using the PCG method with preconditioner denoted by M and with startvector y^0 , we have

$$y^k - y^0 \in \text{span}\{M^{-1}Se^0, \dots, (M^{-1}S)^k e^0\}, \quad e^0 := y - y^0.$$

And thus $\langle y^k - y^0, Me \rangle = 0$ for $k \geq 1$, i.e., $y^k - y^0 \in (Me)^\perp$ for $k \geq 1$. If for the preconditioner we take $M \in \{\hat{M}_\nu, \bar{M}_\nu\}$ then it follows, using (4.8), that

$$y^k \in (\hat{M}_\nu e)^\perp \quad \text{for } k \geq 1, \quad \text{if } y^0 \in (\hat{M}_\nu e)^\perp \tag{4.12}$$

This means that for both preconditioners the iterands remain in the subspace $(\hat{M}_\nu e)^\perp$, if the starting vector y^0 is in this subspace. Since the solution y is also sought in this

subspace (cf. 4.10)) this implies that the errors $e^k := y - y^k$ remain in this subspace if $y^0 \in (\hat{M}_\nu e)^\perp$. Hence, only the spectral condition number of the preconditioned matrix on this subspace is relevant. The results in theorem 4.1 and lemma 4.2 yield that both $\hat{M}_\nu^{-1}S$ and $\bar{M}_\nu^{-1}S$ have optimal (i.e., independent of h and ν) spectral condition numbers on this subspace.

In practice the Uzawa method is not very attractive because one has to solve the A-systems accurately. In this paper we consider the Uzawa method to illustrate the robustness of the multigrid solver and of the preconditioners for the Schur complement (cf. section 5). In practical applications variants of the Uzawa method that are much more efficient are used (cf., for example, [3, 7, 16, 25, 29]). Here we consider a preconditioned MINRES method. For this we consider a symmetric positive definite preconditioner

$$\tilde{K} = \begin{pmatrix} Q_A & 0 \\ 0 & Q_S \end{pmatrix} \quad \text{for } K := \begin{pmatrix} A & B^T \\ B & 0 \end{pmatrix}$$

Define the norm $\|w\|_{\tilde{K}} := \langle \tilde{K}w, w \rangle^{\frac{1}{2}}$ for $w \in \mathbb{R}^{n+m}$. Given a starting vector w^0 with corresponding error $e^0 := w^* - w^0$, then in the preconditioned MINRES method one computes the vector $w^k \in w^0 + \text{span}\{\tilde{K}^{-1}Ke^0, \dots, (\tilde{K}^{-1}K)^k e^0\}$ which minimizes the preconditioned residual $\|\tilde{K}^{-1}K(w^* - w)\|_{\tilde{K}}$ over this subspace. For an efficient implementation of this method we refer to the literature.

If we take $Q_S \in \{\hat{M}_\nu, \bar{M}_\nu\}$, then again we have that the approximations (and errors) of the pressure remain in the subspace $(\hat{M}_\nu e)^\perp$ if the starting approximation y^0 is in this subspace. From the literature (cf. [24, 25]) it is known that the convergence of the preconditioned MINRES method is fast if we have good preconditioners Q_A of A and Q_S of S (on the subspace $(\hat{M}_\nu e)^\perp$). In the numerical experiments in section 5 we take a standard multigrid method for Q_A and $Q_S = \bar{M}_\nu$.

5. Numerical experiments. In this section we present results of a few numerical experiments to illustrate the behaviour of the Uzawa and preconditioned MINRES method applied to the Stokes interface problem. We consider a problem as in (1.1)-(1.3) with

$$\Omega = (0, 1)^3, \quad \Omega_2 = (0, \frac{1}{2})^3$$

For the discretization we start with a uniform tetrahedral grid with $h = \frac{1}{2}$ and we apply regular refinements to this starting triangulation. The resulting triangulations satisfy the conformity condition (3.1). For the finite element discretization we used the LBB stable pair of Hood-Taylor $P_2 - P_1$, i.e. continuous piecewise quadratics for the velocity and continuous piecewise linears for the pressure. We performed computations for the cases $h = 1/16$, $h = 1/32$ and with varying $\varepsilon \in (0, 1]$. Note that for $h = 1/32$ we have approximately $7.5 \cdot 10^5$ velocity unknowns and $3.3 \cdot 10^4$ pressure unknowns ($n \approx 7.5 \cdot 10^5$, $m \approx 3.3 \cdot 10^4$). We consider the linear system as in (4.2) with solution $(x, y) = 0$. We take a fixed arbitrary starting vector (x^0, y^0) , with $y^0 \in (\hat{M}_\nu)^\perp$.

To test the robustness of the Schur complement preconditioning and of the multigrid solver we first consider the Uzawa method (4.9)-(4.11). The linear systems of the form $Ax = r$ that occur in the steps 1,2 and 3 are all solved using a standard

multigrid V-cycle with one pre- and one post-smoothing iteration with a symmetric Gauss-Seidel method. The starting vector is x^0 and the iteration is stopped when for the result after k iterations, x^k , the scaled residual satisfies

$$\frac{\|D^{-1}(Ax^k - r)\|}{\|D^{-1}(Ax^0 - r)\|} \leq 10^{-10}, \quad D := \text{diag}(A) \quad (5.1)$$

Here $\|\cdot\|$ denotes the standard Euclidean norm. The system with the Schur complement in (4.10) is solved using a PCG method with preconditioner \hat{M}_ν . The systems $\hat{M}_\nu y = w$ are solved approximately using a PCG method with preconditioner \bar{M}_ν and starting vector y^0 and accuracy

$$\frac{\|\bar{M}_\nu^{-1}(\hat{M}_\nu y^k - w)\|}{\|\bar{M}_\nu^{-1}(\hat{M}_\nu y^0 - w)\|} \leq 10^{-10} \quad (5.2)$$

The PCG method for the Schur complement system $Sy = c$ has starting vector y^0 and is stopped when

$$\frac{\|\bar{M}_\nu^{-1}(Sy^k - c)\|}{\|\bar{M}_\nu^{-1}(Sy^0 - c)\|} \leq 10^{-6} \quad (5.3)$$

In table 5.1 we present results for different h and ε values. Here #-MG denotes the average number of multigrid iterations needed to satisfy (5.1), #-PCG-M the average number of PCG iterations needed to satisfy (5.2) and #-PCG-S the average number of PCG iterations needed to satisfy (5.3).

These results clearly show the robustness of the multigrid solver for the velocity

TABLE 5.1
Uzawa method, preconditioner \hat{M}_ν

h	1/16				1/32				
	ε	1	10^{-2}	10^{-4}	10^{-6}	1	10^{-2}	10^{-4}	10^{-6}
#-MG		13	13	14	14	14	14	14	14
#-PCG-M		24	25	25	26	24	25	25	25
#-PCG-S		22	29	31	34	21	29	30	34

systems, of the preconditioner \hat{M}_ν for S and of the preconditioner \bar{M}_ν for \hat{M}_ν with respect to variation of h and of ε . We now consider the effect of using the lumped mass matrix \bar{M}_ν instead of \hat{M}_ν as a preconditioner for the Schur complement. In the PCG method we use a stopping criterion as in (5.3). Note that in this PCG method the preconditioner is now a diagonal matrix. In table 5.2 we present results for different h and ε values. As expected, the lumped mass matrix \bar{M}_ν is a robust preconditioner

TABLE 5.2
Uzawa method, preconditioner \bar{M}_ν

h	1/16				1/32				
	ε	1	10^{-2}	10^{-4}	10^{-6}	1	10^{-2}	10^{-4}	10^{-6}
#-PCG-S		40	48	48	58	39	50	52	59

for the Schur complement S. In the final experiment we consider the preconditioned MINRES method. For the preconditioner Q_A we take *one* iteration of the multigrid

method described above and we take the lumped mass matrix \bar{M}_ν as preconditioner for the Schur complement. In table 5.3 we show the number of iterations k (denoted by #-PMINRES), such that

$$\left\| \tilde{K}^{-1}K \begin{pmatrix} x^k \\ y^k \end{pmatrix} \right\|_{\tilde{K}} \leq 10^{-6} \left\| \tilde{K}^{-1}K \begin{pmatrix} x^0 \\ y^0 \end{pmatrix} \right\|_{\tilde{K}}$$

(recall that the right handside is 0). Note that for $h = 1/32$ one needs less iterations

TABLE 5.3
MINRES with preconditioner \bar{M}_ν

h	1/16				1/32			
ε	1	10^{-2}	10^{-4}	10^{-6}	1	10^{-2}	10^{-4}	10^{-6}
#-PMINRES	62	68	98	157	50	58	85	116

than for $h = 1/16$.

6. Comments on closely related problems. We comment on extensions of the analysis two other Stokes interface problems that are very similar to problem (1.6).

Firstly, we consider the case with the full velocity tensor $D(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$ instead of $\nabla \mathbf{u}$ in (1.1). The corresponding weak formulation is as follows (cf. (1.6)): given $\mathbf{f} \in \mathbf{V}'$ find $\{\mathbf{u}, p\} \in \mathbf{V} \times M$ such that

$$\begin{cases} \int_{\Omega} \nu D(\mathbf{u}) : D(\mathbf{v}) \, dx - (\operatorname{div} \mathbf{v}, p) = (\mathbf{f}, \mathbf{v}) & \text{for } \mathbf{v} \in \mathbf{V}, \\ (\operatorname{div} \mathbf{u}, q) = 0 & \text{for } q \in M. \end{cases} \quad (6.1)$$

Note that

$$D(\mathbf{u}) : D(\mathbf{v}) = \sum_{i,j=1}^d D(\mathbf{u})_{i,j} D(\mathbf{v})_{i,j} = \operatorname{tr}(D(\mathbf{u})D(\mathbf{v}))$$

For $\mathbf{u} \in \mathbf{V}$ we introduce the notation $\|\mathbf{u}\|_{\mathbf{V}_D}^2 := \int_{\Omega} \nu \operatorname{tr}(D(\mathbf{u})^2) \, dx$. In the following lemma we show that this functional defines a norm on \mathbf{V} that is uniformly w.r.t. ν equivalent to $\|\cdot\|_{\mathbf{V}}$.

LEMMA 6.1. *Assume that (2.13) or (2.14) holds. Then there exists a constant $c > 0$ independent of ν such that*

$$c\|\mathbf{v}\|_{\mathbf{V}} \leq \|\mathbf{v}\|_{\mathbf{V}_D} \leq \|\mathbf{v}\|_{\mathbf{V}} \quad \text{for all } \mathbf{v} \in \mathbf{V} \quad (6.2)$$

Proof. First note that

$$\operatorname{tr}(D(\mathbf{v})^2) = \frac{1}{4} \sum_{i,j=1}^d \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right)^2 \leq \sum_{i,j=1}^d \left(\frac{\partial u_i}{\partial x_j} \right)^2$$

This yields

$$\|\mathbf{v}\|_{\mathbf{V}_D}^2 = \int_{\Omega} \nu \operatorname{tr}(D(\mathbf{v})^2) \, dx \leq (\nu \nabla \mathbf{v}, \nabla \mathbf{v}) = \|\mathbf{v}\|_{\mathbf{V}}^2$$

and thus the second inequality in (6.2) holds. To prove the first inequality in (6.2) we use Korn's inequality, cf. [13, 18]. Let γ be a part of $\partial\Omega$ with nonzero measure, then there exists a constant C such that

$$\|\nabla\mathbf{v}\|_{\Omega}^2 := \|\nabla\mathbf{v}\|_{L^2(\Omega)}^2 \leq C \int_{\Omega} \text{tr}(D(\mathbf{v})^2) dx$$

for all $\mathbf{v} \in H^1(\Omega)^d$ with $\mathbf{v}|_{\gamma} = 0$. Assume that (2.13) holds. Korn's inequality holds in each of the two subdomains and thus we get

$$\|\nabla\mathbf{v}\|_{\Omega_i}^2 \leq C_i \int_{\Omega_i} \text{tr}(D(\mathbf{v})^2) dx \quad \text{for } i = 1, 2 \quad (6.3)$$

Hence

$$\begin{aligned} \|\mathbf{v}\|_{\mathbf{V}}^2 &= \|\nabla\mathbf{v}\|_{\Omega_1}^2 + \varepsilon \|\nabla\mathbf{v}\|_{\Omega_2}^2 \\ &\leq \max\{C_1, C_2\} \int_{\Omega} \nu \text{tr}(D(\mathbf{v})^2) dx = \max\{C_1, C_2\} \|\mathbf{v}\|_{\mathbf{V}_D}^2 \end{aligned}$$

which yields the first inequality in (6.2). Now assume that (2.14) holds. We then have the result in (6.3) for $i = 1$. Using a trace theorem we get

$$\|\nabla\mathbf{v}\|_{\Omega_2}^2 \leq c \|\mathbf{v}|_{\Gamma}\|_{H^{\frac{1}{2}}(\Gamma)}^2 \leq c \|\nabla\mathbf{v}\|_{\Omega_1}^2 \leq c \int_{\Omega_1} \text{tr}(D(\mathbf{v})^2) dx$$

Using the assumption that ε is bounded we obtain

$$\|\mathbf{v}\|_{\mathbf{V}}^2 = \|\nabla\mathbf{v}\|_{\Omega_1}^2 + \varepsilon \|\nabla\mathbf{v}\|_{\Omega_2}^2 \leq C \int_{\Omega_1} \text{tr}(D(\mathbf{v})^2) dx \leq C \|\mathbf{v}\|_{\mathbf{V}_D}^2$$

and thus the first inequality in (6.2) holds. \square

Assume that (2.13) or (2.14) holds. Due to this lemma we can conclude that the continuity result in lemma 2.1 and the infsup properties in theorem 2.2 and theorem 3.3 also hold with $\|\cdot\|_{\mathbf{V}_D}$ instead of $\|\cdot\|_{\mathbf{V}}$. Thus our analysis also applies to the variational problem (6.1).

We note that the assumptions (2.13), (2.14) are not essential. The upper bound in (6.2) holds without these assumptions, and thus the infsup properties in the norm $\|\cdot\|_{\mathbf{V}_D}$, too. A continuity result as lemma 2.1 with $\|\cdot\|_{\mathbf{V}_D}$ instead of $\|\cdot\|_{\mathbf{V}}$ can be proved without assuming (2.13), (2.14).

The second comment is related to the choice of the pressure space. Instead of the pressure space M introduced in (1.5) one may want to use the standard space

$$L_0^2(\Omega) = \{p \in L^2(\Omega) \mid (p, 1) = 0\}.$$

and consider the variational formulation of the interface problem (1.1)-(1.3) in the space $\mathbf{V} \times L_0^2(\Omega)$. It turns out that the analysis then requires different norms to obtain optimal estimates. These optimal estimates are ν -independent for the continuous problem. For the corresponding Galerkin discrete problem, however, opposite to the results in section 3 we now observe a ν -dependence in the estimates. These results are presented in [19]. The main ideas of the analysis in [19] are the same as in the sections 2 and 3. However, some further technicalities, like a mesh-dependent norm in the pressure space, are needed.

Acknowledgements. The authors thank J. Peters for providing the results of the numerical experiments

REFERENCES

- [1] Babuska, I. (1970): The finite element method for elliptic problems with discontinuous coefficients, *Computing*, **5**, 2-7–213.
- [2] Babuska, I., Caloz, C., Osborn, J. (1994): Special finite element methods for a class of second order elliptic problems with rough coefficients, *SIAM J. Numer. Anal.*, **31**, 945–981.
- [3] Bank, R.E., Welfert, B.D., and Yserentant, H. (1990): A class of iterative methods for solving saddle point problems, *Numer. Math.*, **56**, 645–666.
- [4] Bernardi, C., Verfürth, R. (2000): Adaptive finite element methods for elliptic equations with non-smooth coefficients, *Numer. Math.*, **85**, 579–608
- [5] Brackbill, J.U., Kothe, D.B., Zemach, C. (1992): A continuum method for modeling surface tension, *J. Comput. Phys.*, **100**, 335–354.
- [6] Bramble, J.H. and King, J.T. (1996): A finite element method for interface problems in domains with smooth boundaries and interfaces, *Advances in Comp. Math.*, **6**, 109–138.
- [7] Bramble, J.H., Pasciak, J.E., and Vassilev, A.T. (1997): Analysis of the inexact Uzawa algorithm for saddle point problems. *SIAM J. Numer. Anal.*, **34**, 1072–1092.
- [8] Bramble, J.H., *Multigrid Methods*, Pitman Research Notes in Mathematics, Vol. 294, John Wiley, New York, 1993.
- [9] Bramble, J.H., Pasciak, J.E., Wang, J., Xu, J. (1991): Convergence estimates for multigrid algorithms without regularity assumptions. *Math. Comp.*, **57**, 23–45.
- [10] Brezzi, F. and Fortin, M., *Mixed and Hybrid Finite Element Methods*, Springer, New York, 1991.
- [11] Ciarlet, P.G., *The Finit Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [12] Ciarlet, P.G.: Basic Error Estimates for Elliptic Problems. *Handbook of Numerical Analysis*, Vol. II, P.G. Ciarlet and J.L. Lions, eds., Elsevier Science, 1991, 17–351.
- [13] Dautray, R., Lions, J.-L., *Mathematical Analysis and Numerical Methods for Science and Technology, Vol. 2: Functional and Variational Methods*, Springer, Berlin, 1988.
- [14] Chang, Y.C., Hou, T.Y., Merriman, B., Osher, S. (1996): Eulerian capturing methods based on a level set formulation for incompressible fluid interfaces, *J. Comput. Phys.*, **124**, 449–464.
- [15] Chen, Z. and Zou, J. (1998): Finite element method and its convergence for elliptic and parabolic interfac problems. *Numer. Math.* **79**, 175–202
- [16] Elman, H.C. and Golub, G.H. (1994): Inexact and preconditioned Uzawa algorithms for saddle point problems, *SIAM J. Numer. Anal.*, **31**, 1645–1661.
- [17] Girault, V., and Raviart, P.-A., *Finite Element Methods for Navier-Stokes Equations*, Springer, Berlin, 1986.
- [18] Kondratiev, V.A., Oleinik, O.A. (1988): Boundary-value problems for the systems of elasticity in unbounded domains. Korn’s inequalities, *Russian Math. Surveys*, **43**, 65–119
- [19] Olshanski, M.A. Reusken, A. (2004): A Stokes Interface Problem: Stability, Finite Element Analysis and a Robust Solver, *Proc. of European Congress on Computational Methods in Applied Sciences and Engineering, ECCOMAS 2004* (Eds. P. Neittaanmäki, et al.), 2004
- [20] Osher, S., Fedkiw, R.P. (2001): Level set methods: An overview and some recent results, *J. Comput. Phys.*, **169**, 463–502.
- [21] Petzoldt, M. (2002): A posteriori error estimators for elliptic equations with discontinuous diffusion coefficients. *Advances in Computational Mathematics*, **16**, 47–75.
- [22] Plum, M. and Wieners, C. (2003): Optimal a priori estimates for interface problems. *Numer. Math.* **95**, 735–759
- [23] Powell, C., Silvester, D. (2003): Black-box preconditioning for mixed formulation of self-adjoint elliptic PDEs. In: *Challenges in Scientific Computing - CISC 2002*, ed. E. Bänsch. Lecture Notes in Computational Science and Engineering, Vol. 35, 268–285.
- [24] Rusten, T. and Winther, R. (1992): A preconditioned iterative method for saddlepoint problems. *SIAM J. Matrix Anal. Appl.*, **13**, 887–904.
- [25] Silvester, D. Wathen, A (1994): Fast iterative solution of stabilised stokes systems. part II: using general block preconditioners. *SIAM J. Numer. Anal.*, **31**, 1352–1367.
- [26] Tornberg, A.-K., Engquist, B. (2000): A finite element based level-set method for multiphase flow applications, *Comput. Visual. Sci.*, **3**, 93–101.
- [27] Wathen, A. (1987): Realistic eigenvalues bounds for the Galerkin mass matrix. *IMA J. Numer. Anal.*, **7**, 449–457.

- [28] Yserentant, H. (1990): Two preconditioners based on the multi-level splitting of finite element spaces. *Numer. Math.*, **58**, 163–184.
- [29] Zulehner, W. (2001): Analysis of iterative methods for saddle point problems: A unified approach, *Math. Comp.*, **71**, 479–505.